

DATE

The Honorable E. Scott Pruitt  
Administrator  
U.S. Environmental Protection Agency  
1200 Pennsylvania Avenue, N.W.  
Washington, DC 20460

Subject: Review of EPA's Draft Assessment entitled *Toxicological Review of Hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX)* (September 2016)

Dear Administrator Pruitt:

The EPA's National Center for Environmental Assessment (NCEA) requested that the Science Advisory Board (SAB) review the draft assessment, entitled *Draft Toxicological Review of Hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX)*. The draft assessment consists of a review of available scientific literature on the toxicity of RDX. The SAB was asked to comment on the scientific soundness of the hazard and dose-response assessment of RDX-induced cancer and noncancer health effects. In response to EPA's request, the SAB convened a panel consisting of members of the SAB Chemical Assessment Advisory Committee (CAAC) augmented with subject matter experts to conduct the review.

The SAB finds the draft assessment to be comprehensive and generally well-written. The enclosed report provides the SAB's consensus advice and recommendations. This letter briefly conveys the major findings.

The draft assessment evaluates and modifies available physiologically-based pharmacokinetic (PBPK) models in the literature. The SAB finds the revised rat and human PBPK models to be a distinct improvement over the original approach, and these changes adequately represent RDX toxicokinetics. The application of revised PBPK models in the assessment for the calculation of human equivalent doses (HEDs) for the points of departure (PODs) for neurotoxicity and other noncancer endpoints is scientifically supported. For the hazard identification and dose-response assessment of noncancer endpoints, the SAB agrees that neurotoxicity, including seizures or convulsions, is a human hazard of RDX exposure. However, convulsions in rodents only provide a limited spectrum of the potential human hazard, since convulsive or non-convulsive seizures, epileptiform discharges, reduction in seizure threshold, subchronic sensitization, and neuronal damage can all be part of the spectrum of RDX's nervous system hazards. Thus, further evaluation or explanation should be provided in the draft assessment for these potential endpoints. The SAB agrees that RDX-induced convulsions arise primarily through a mode of action involving RDX-induced gamma-amino butyric acid type A (GABA<sub>A</sub>) receptor (GABA<sub>A</sub>R) blockade. The SAB also agrees with the characterization of convulsions as a severe endpoint, and concludes that its potential relationship to mortality is clearly described. However, the SAB recommends that EPA revisit the benchmark response (BMR), and at a minimum,

1 improve their justification for using a BMR of 1% for deriving the lower bound on the  
2 benchmark dose (BMDL) as the point of departure (POD) from Crouse et al. (2006). Given that a  
3 BMR of 1% corresponds to a response that is a factor of 15 below the lowest observed response  
4 data, the SAB considers the use of BMR of 5% based on the Crouse study to be more consistent  
5 with the observed response at the Lowest-Observed-Adverse-Effect-Level (LOAEL) of 15%,  
6 and not so far below the observable data. Thus, EPA should consider use of a 5% BMR while  
7 addressing the uncertainty of a frank effect with uncertainty factors, or at least provide a more  
8 thorough justification for its choice of a 1% BMR.

9  
10 With respect to the application of uncertainty factors (UF) to the PODs, the SAB supports the  
11 application of an interspecies UF of 3 to account for the toxicodynamic and residual  
12 toxicokinetic uncertainty in extrapolation from animal to human not accounted for by the  
13 toxicokinetic modeling. In addition, the SAB agrees with the LOAEL to No-Observed-Adverse-  
14 Effect-Level (NOAEL) UF of 1, and the UF of 10 to account for intra-human variability.  
15 However, the SAB has concerns about the use of a subchronic to chronic UF (UF<sub>S</sub>) of 1.  
16 Observations in an *in vitro* assay for GABA activity found that the effects of RDX were not  
17 reversible following compound wash out (Williams et al. 2011). As such, it is possible that  
18 repeated exposures to RDX might have cumulative effects on GABAergic neurotransmission.  
19 The SAB recommends that EPA reconsider the UF for subchronic to chronic extrapolation, and  
20 that at a minimum, provide stronger justification for a UF<sub>S</sub> of 1. In addition, the SAB questions  
21 the application of a database uncertainty factor (UF<sub>D</sub>) of 3, and suggests EPA consider applying a  
22 UF<sub>D</sub> of 10 to account for data gaps for developmental neurotoxicity, lack of incidence data for  
23 less severe nervous system effects, and proximity of the dose that induces convulsions with the  
24 dose that induces mortality. A composite UF of 300 should be considered instead of the UF of  
25 100 proposed in the draft assessment.

26  
27 The SAB concludes that the derived reference dose (RfD) for nervous system effects is not  
28 scientifically supported as it did not capture all of the potential adverse nervous system outcomes  
29 or their severity, and it does not account for many database uncertainties. The SAB recommends  
30 the EPA use the dose-response data of the Crouse et al. (2006) study as the primary basis for the  
31 derivation of an RfD for neurotoxicity. In particular, the SAB concludes that the functional  
32 observation battery (FOB) data presented in the Crouse study were rudimentary and not  
33 sufficiently sensitive to detect neurobehavioral consequences produced by chronic/subchronic  
34 doses of RDX over prolonged periods, especially during pregnancy. Moreover, tests directed at  
35 detecting subtle developmental neurotoxicity during the perinatal-weaning period have not been  
36 conducted. These concerns are especially compelling because of more recent peer-reviewed  
37 published data indicating that sub-convulsive doses of either bicuculline (which has a similar  
38 mechanism of action to RDX) or domoic acid (which has agonist activity on glutamate  
39 transmission) have been shown to cause developmental and behavioral impairments at doses  
40 below those that cause convulsions. Thus, the SAB concludes that there remains significant  
41 uncertainty about the developmental neurotoxicity of RDX.

42  
43 The SAB agrees that kidney and other urogenital system toxicity are a potential human hazard of  
44 RDX exposure. However, the SAB disagrees with the selection of suppurative prostatitis as the  
45 “surrogate marker” to represent this hazard, and recommends that EPA considers suppurative  
46 prostatitis as a separate effect. As such, separate organ/system-specific RfDs should be derived

1 for the kidney and urogenital system, based on findings of renal papillary necrosis and associated  
2 renal inflammation, and for suppurative prostatitis.

3  
4 The SAB disagrees with the conclusion that male reproductive effects are a human hazard  
5 associated with RDX exposure. The SAB concludes that RDX does not pose a risk of induction  
6 of structural malformations during human fetal development based on animal data. Additionally,  
7 the SAB agrees that conclusions cannot be drawn regarding other forms of developmental  
8 toxicity, which were only seen at maternally toxic dose levels. The SAB also notes that potential  
9 neurodevelopmental toxicity based on the reported mechanism of RDX inhibition of GABAergic  
10 neurons, and the findings that RDX is present in the brain of offspring rats and in the milk from  
11 dams treated with RDX during gestation, were not adequately discussed in the draft assessment.  
12 The SAB concludes the proposed RfD for reproductive system effects in the draft assessment is  
13 not scientifically supported.

14  
15 With regard to cancer effects, the SAB agrees that “suggestive evidence of carcinogenic  
16 potential” is the most appropriate cancer hazard descriptor for RDX, in accordance with EPA’s  
17 *Guidelines for Carcinogen Risk Assessment*; and that this descriptor applies to all routes of  
18 exposure. The SAB also agrees with the agency’s rationale for a quantitative cancer dose-  
19 response analysis for RDX and the use of the linear low-dose extrapolation approach, since the  
20 mode of action for cancer is unknown. However, the SAB finds that the calculations of the  
21 PODs and oral slope factor were not clearly described, and questions whether these are  
22 scientifically supported. The SAB recognizes the Agency’s preference for using the multistage  
23 model for cancer dose-response modeling. However, the SAB identifies a number of concerns  
24 with the data used to derive the cancer POD, the rationale for restricting modeling to the  
25 multistage model to derive the POD, and the conditions under which the agency’s MS-COMBO  
26 modeling methodology provides a valid POD and cancer slope factor estimate. The SAB also  
27 makes multiple suggestions on how the discussion on the derivation of the oral slope factor can  
28 be improved.

29  
30 With regard to dose-response analysis of noncancer effects, the SAB agrees that the overall RfD  
31 should be based on nervous system effects. The SAB recommends that the EPA use the dose-  
32 response data of the Crouse et al. (2006) study as the primary basis for the derivation of an RfD.  
33 An RfD derived from the NOAEL of Cholakis et al. (1980) can be used for comparison.

34  
35 The SAB appreciates this opportunity to review EPA’s *Draft Toxicological Review of RDX* and  
36 looks forward to the EPA’s response to these recommendations.

37  
38 Sincerely,

39  
40  
41  
42 Dr. Peter S. Thorne, Chair  
43 EPA Science Advisory Board

44 Dr. Kenneth S. Ramos, Chair  
45 SAB CAAC Augmented for the Review of the  
46 Draft IRIS RDX Assessment

47 Enclosure

**NOTICE**

This report has been written as part of the activities of the EPA Science Advisory Board, a public advisory committee providing extramural scientific information and advice to the Administrator and other officials of the Environmental Protection Agency. The Board is structured to provide balanced, expert assessment of scientific matters related to problems facing the Agency. This report has not been reviewed for approval by the Agency and, hence, the contents of this report do not represent the views and policies of the Environmental Protection Agency, nor of other agencies in the Executive Branch of the Federal government, nor does mention of trade names or commercial products constitute a recommendation for use. Reports of the EPA Science Advisory Board are posted on the EPA website at <http://www.epa.gov/sab>.

**U.S. Environmental Protection Agency**  
**Science Advisory Board**  
**Chemical Assessment Advisory Committee Augmented for the**  
**Review of Draft IRIS RDX Assessment**

**CHAIR**

**Dr. Kenneth Ramos**, Associate Vice-President of Precision Health Sciences and Professor of Medicine, Arizona Health Sciences Center, University of Arizona, Tucson, AZ

**MEMBERS**

**Dr. Hugh A. Barton**, Associate Research Fellow, Pharmacokinetics, Dynamics, and Metabolism, Pfizer Inc., Groton, CT

**Dr. Maarten C. Bosland**, Professor of Pathology, College of Medicine, University of Illinois at Chicago, Chicago, IL

**Dr. Mary Boudreau**, Research Toxicologist, Division of Biochemical Toxicology, National Center for Toxicological Research, U.S. Food and Drug Administration, Jefferson, AR

**Dr. James V. Bruckner**, Professor, Department of Pharmacology & Toxicology, College of Pharmacy, University of Georgia, Athens, GA

**Dr. George Cobb**, Professor, Environmental Science, College of Arts and Sciences, Baylor University, Waco, TX

**Dr. David Eastmond**, Professor and Chair, Department of Cell Biology and Neuroscience, Toxicology Graduate Program, University of California at Riverside, Riverside, CA

**Dr. Joanne English**, Independent Consultant, Menlo Park, CA

**Dr. Alan Hoberman**, Toxicologist, Research, Charles River Laboratories, Inc., Horsham, PA

**Dr. Jacqueline Hughes-Oliver**, Professor, Statistics Department, North Carolina State University, Raleigh, NC

**Dr. Susan Laffan**, Safety Assessment, GlaxoSmithKline, King of Prussia, PA

**Dr. Lawrence Lash**, Professor, Department of Pharmacology, Wayne State University School of Medicine, Wayne State University, Detroit, MI

**Dr. Stephen Lasley**, Professor of Pharmacology and Assistant Head, Cancer Biology & Pharmacology, College of Medicine, University of Illinois at Chicago, Peoria, IL

**Dr. Melanie Marty**, Adjunct Professor, Environmental Toxicology, University of California at Davis, Davis, CA

1  
2 **Dr. Marvin Meistrich**, Professor, Experimental Radiation Oncology, M.D. Anderson Cancer  
3 Center, University of Texas, Houston, TX  
4

5 **Dr. Marilyn Morris**, Professor of Pharmaceutical Sciences, School of Pharmacy and  
6 Pharmaceutical Sciences, University at Buffalo, State University of New York, Buffalo, NY  
7

8 **Dr. Victoria Persky**, Professor, Epidemiology & Biostatistics Program, School of Public Health,  
9 University of Illinois at Chicago, Chicago, IL  
10

11 **Dr. Isaac Pessah**, Professor, Molecular Biosciences, School of Veterinary Medicine, University  
12 of California at Davis, Davis, CA  
13

14 **Dr. Kenneth M. Portier**, Vice President, Department of Statistics & Evaluation Center,  
15 American Cancer Society, Atlanta, GA  
16

17 **Dr Samba Reddy**, Professor, Neuroscience and Experimental Therapeutics, College of  
18 Medicine, Texas A&M University, Bryan, TX  
19

20 **Dr. Stephen M. Roberts**, Professor, Center for Environmental and Human Toxicology,  
21 University of Florida, Gainesville, FL  
22

23 **Dr. Thomas Rosol**, Professor, Veterinary Biosciences, College of Veterinary Medicine, Ohio  
24 State University, Columbus, OH  
25

26 **Dr. Alan Stern**, Chief, Bureau for Risk Analysis, Division of Science, Research and  
27 Environmental Health, New Jersey Department of Environmental Protection, Trenton, NJ  
28

29 **Dr. Robert Turesky**, Professor, Masonic Cancer Center and Department of Medicinal  
30 Chemistry, College of Pharmacy, University of Minnesota, Minneapolis, MN  
31

32  
33 **SCIENCE ADVISORY BOARD STAFF**  
34

35 **Dr. Diana Wong**, Designated Federal Officer, U.S. Environmental Protection Agency, Science  
36 Advisory Board (1400R), 1200 Pennsylvania Avenue, NW, Washington, DC  
37  
38  
39

**U.S. Environmental Protection Agency  
Science Advisory Board  
BOARD**

**CHAIR**

**Dr. Peter S. Thorne**, Professor and Head, Department of Occupational & Environmental Health, College of Public Health, University of Iowa, Iowa City, IA

**MEMBERS**

**Dr. Joseph Arvai**, Max McGraw Professor of Sustainable Enterprise and Director, Erb Institute, School of Natural Resources & Environment, University of Michigan, Ann Arbor, MI

**Dr. Deborah Hall Bennett**, Professor and Interim Chief, Environmental and Occupational Health Division, Department of Public Health Sciences, School of Medicine, University of California, Davis, Davis, CA

**Dr. Kiros T. Berhane**, Professor, Preventive Medicine, Keck School of Medicine, University of Southern California, Los Angeles, CA

**Dr. Sylvie M. Brouder**, Professor and Wickersham Chair of Excellence in Agricultural Research, Department of Agronomy, Purdue University, West Lafayette, IN

**Dr. Joel G. Burken**, Curator's Professor and Chair, Civil, Architectural, and Environmental Engineering, College of Engineering and Computing, Missouri University of Science and Technology, Rolla, MO, United States

**Dr. Janice E. Chambers**, William L. Giles Distinguished Professor and Director, Center for Environmental Health and Sciences, College of Veterinary Medicine, Mississippi State University, Starkville, MS

**Dr. Alison C. Cullen**, Professor, Daniel J. Evans School of Public Policy and Governance, University of Washington, Seattle, WA

**Dr. Ana V. Diez Roux**, Dean, School of Public Health, Drexel University, Philadelphia, PA  
Also Member: CASAC

**Dr. Otto C. Doering III**, Professor, Department of Agricultural Economics, Purdue University, W. Lafayette, IN

**Dr. Joel J. Ducoste**, Professor, Department of Civil, Construction, and Environmental Engineering, College of Engineering, North Carolina State University, Raleigh, NC

**Dr. Susan P. Felter**, Research Fellow, Global Product Stewardship, Procter & Gamble, Mason, OH

**Dr. R. William Field**, Professor, Department of Occupational and Environmental Health and  
Department of Epidemiology, College of Public Health, University of Iowa, Iowa City, IA

**Dr. H. Christopher Frey**, Glenn E. Futrell Distinguished University Professor, Department of  
Civil, Construction and Environmental Engineering, College of Engineering, North Carolina  
State University, Raleigh, NC

**Dr. Joseph A. Gardella**, SUNY Distinguished Professor and John and Frances Larkin Professor  
of Chemistry, Department of Chemistry, College of Arts and Sciences, University at Buffalo,  
Buffalo, NY

**Dr. Steven P. Hamburg**, Chief Scientist, Environmental Defense Fund, Boston, MA

**Dr. Cynthia M. Harris**, Director and Professor, Institute of Public Health, Florida A&M  
University, Tallahassee, FL

**Dr. Robert J. Johnston**, Director of the George Perkins Marsh Institute and Professor,  
Department of Economics, Clark University, Worcester, MA

**Dr. Kimberly L. Jones**, Professor and Chair, Department of Civil and Environmental  
Engineering, Howard University, Washington, DC

**Dr. Catherine J. Karr**, Associate Professor - Pediatrics and Environmental and Occupational  
Health Sciences and Director - NW Pediatric Environmental Health Specialty Unit, University of  
Washington, Seattle, WA

**Dr. Madhu Khanna**, ACES Distinguished Professor in Environmental Economics, Director of  
Graduate Admissions and Associate Director, Institute of Sustainability, Energy, and  
Environment, Department of Agricultural and Consumer Economics, University of Illinois at  
Urbana-Champaign, Urbana, IL

**Dr. Francine Laden**, Professor of Environmental Epidemiology, Associate Chair Environmental  
Health and Director of Exposure, Departments of Environmental Health and Epidemiology ,  
Harvard T.H. Chan School of Public Health, Boston, MA

**Dr. Robert E. Mace**, Deputy Executive Administrator, Water Science & Conservation, Texas  
Water Development Board, Austin, TX

**Dr. Clyde F. Martin**, Horn Professor of Mathematics, Emeritus, Department of Mathematics  
and Statistics, Texas Tech University, Crofton, MD

**Dr. Sue Marty**, Senior Toxicology Leader, Toxicology & Environmental Research, The Dow  
Chemical Company, Midland, MI

**Dr. Denise Mauzerall**, Professor, Woodrow Wilson School of Public and International Affairs,



and Department of Civil and Environmental Engineering, Princeton University, Princeton, NJ

**Dr. Kristina D. Mena**, Associate Professor, Epidemiology, Human Genetics and Environmental Sciences, School of Public Health, University of Texas Health Science Center at Houston, El Paso, TX

**Dr. Surabi Menon**, Director of Research, ClimateWorks Foundation, San Francisco, CA

**Dr. Kari Nadeau**, Naddisy Family Foundation Professor of Medicine, Director, FARE Center of Excellence at Stanford University, and Sean N. Parker Center for Allergy and Asthma Research at, Stanford University School of Medicine, Stanford, CA

**Dr. James Opaluch**, Professor and Chair, Department of Environmental and Natural Resource Economics, College of the Environment and Life Sciences, University of Rhode Island, Kingston, RI

**Dr. Thomas F. Parkerton**, Senior Environmental Associate, Toxicology & Environmental Science Division, ExxonMobil Biomedical Science, Houston, TX

**Mr. Richard L. Poirot**, Independent Consultant, Independent Consultant, Burlington, VT

**Dr. Kenneth M. Portier**, Vice President, Department of Statistics & Evaluation Center, American Cancer Society, Atlanta, GA

**Dr. Kenneth Ramos**, Associate Vice-President of Precision Health Sciences and Professor of Medicine, Arizona Health Sciences Center, University of Arizona, Tucson, AZ

**Dr. David B. Richardson**, Associate Professor, Department of Epidemiology, School of Public Health, University of North Carolina, Chapel Hill, NC

**Dr. Tara L. Sabo-Attwood**, Associate Professor and Chair, Department of Environmental and Global Health, College of Public Health and Health Professionals, University of Florida, Gainesville, FL

**Dr. William Schlesinger**, President Emeritus, Cary Institute of Ecosystem Studies, Millbrook, NY

**Dr. Gina Solomon**, Deputy Secretary for Science and Health, Office of the Secretary, California Environmental Protection Agency, Sacramento, CA

**Dr. Daniel O. Stram**, Professor, Department of Preventive Medicine, Division of Biostatistics, University of Southern California, Los Angeles, CA

**Dr. Jay Turner**, Associate Professor and Vice Dean for Education, Department of Energy, Environmental and Chemical Engineering, School of Engineering & Applied Science, Washington University, St. Louis, MO

**Dr. Edwin van Wijngaarden**, Associate Professor, Department of Public Health Sciences,  
School of Medicine and Dentistry, University of Rochester, Rochester, NY

**Dr. Jeanne M. VanBriesen**, Duquesne Light Company Professor of Civil and Environmental  
Engineering, and Director, Center for Water Quality in Urban Environmental Systems (Water-  
QUEST), Department of Civil and Environmental Engineering, Carnegie Mellon University,  
Pittsburgh, PA

**Dr. Elke Weber**, Gerhard R. Andlinger Professor in Energy and the Environment, Professor of  
Psychology and Public Affairs, Woodrow Wilson School of Public and International Affairs,  
Princeton University, Princeton, NJ

**Dr. Charles Werth**, Professor and Bettie Margaret Smith Chair in Environmental Health  
Engineering, Department of Civil, Architectural and Environmental Engineering, Cockrell  
School of Engineering, University of Texas at Austin, Austin, TX

**Dr. Peter J. Wilcoxon**, Laura J. and L. Douglas Meredith Professor for Teaching Excellence,  
Director, Center for Environmental Policy and Administration, The Maxwell School, Syracuse  
University, Syracuse, NY

**Dr. Robyn S. Wilson**, Associate Professor, School of Environment and Natural Resources, Ohio  
State University, Columbus, OH

#### **SCIENCE ADVISORY BOARD STAFF**

**Mr. Thomas Carpenter**, Designated Federal Officer, U.S. Environmental Protection Agency,  
Science Advisory Board (1400R), 1200 Pennsylvania Avenue, NW, Washington, DC

## TABLE OF CONTENTS

<b>ABBREVIATIONS AND ACRONYMS.....</b>	<b>12</b>
<b>1. EXECUTIVE SUMMARY.....</b>	<b>14</b>
<b>2. INTRODUCTION.....</b>	<b>20</b>
<b>3. RESPONSES TO EPA’S CHARGE QUESTIONS .....</b>	<b>21</b>
<b>3.1. LITERATURE SEARCH/STUDY SELECTION AND EVALUATION.....</b>	<b>21</b>
<b>3.2. TOXICOKINETIC MODELING .....</b>	<b>25</b>
3.2.1. Model Evaluation.....	25
3.2.2. Selection of Dose Metric .....	27
3.2.3. Intrahuman Variation .....	28
<b>3.3. HAZARD IDENTIFICATION AND DOSE-RESPONSE ASSESSMENT .....</b>	<b>29</b>
3.3.1. Nervous System Effects .....	29
3.3.2. Kidney and Other Urogenital System Effects.....	45
3.3.3. Developmental and Reproductive System Effects.....	52
3.3.4. OTHER NONCANCER HAZARDS .....	59
3.3.5. Cancer .....	62
<b>3.4. DOSE-RESPONSE ANALYSIS.....</b>	<b>70</b>
3.4.1. Oral Reference Dose for Effects other than Cancer.....	70
3.4.2. Inhalation Reference Concentration for Effects other than Cancer .....	74
3.4.3. Oral Slope Factor for Cancer .....	75
3.4.4. Inhalation Unit Risk for Cancer.....	77
<b>3.5. EXECUTIVE SUMMARY.....</b>	<b>77</b>
<b>REFERENCES.....</b>	<b>79</b>
<b>APPENDIX A: EPA’S CHARGE QUESTIONS .....</b>	<b>A-1</b>
<b>APPENDIX B: EDITORIAL COMMENTS .....</b>	<b>B-1</b>
<b>APPENDIX C: SUGGESTIONS ON THE FORMAT FOR EPA’S CHARGE QUESTIONS .....</b>	<b>C-1</b>

## ABBREVIATIONS AND ACRONYMS

1	
2	
3	AIC Akaike Information Criteria
4	ATSDR Agency for Toxic Substances and Disease Registry
5	AUC Area Under the Curve
6	BDNF Brain-Derived Neurotrophic Factor
7	BLA Basolateral Amygdala
8	BMC Benchmark Concentration
9	BMCL Lower 95% Confidence Limit of the Benchmark Concentration
10	BMD Benchmark Dose
11	BMDL Lower 95% Confidence Limit of the Benchmark Dose
12	BMR Benchmark Response
13	BW Body Weight
14	CAAC Chemical Assessment Advisory Committee
15	CI Confidence Interval
16	EPA Environmental Protection Agency
17	GABA Gamma-Amino Butyric Acid
18	GABA <sub>A</sub> R Gamma-Amino Butyric Acid Type A Receptor
19	GABAergic Pertaining to or affecting Gamma-Amino Butyric Acid
20	HED Human Equivalent Dose
21	HERO Health and Environmental Research Online
22	HMX Octahydro-1,3,5,7-tetranitro-1,3,5,7-tetrazocine
23	IARC International Agency for Research on Cancer
24	ILSI International Life Sciences Institute
25	IPSPs Inhibitory Postsynaptic Potentials
26	IRIS Integrated Risk Information System
27	IUR Inhalation Unit Risk
28	K <sub>i</sub> Inhibition Constant
29	LD Lethal Dose
30	LOAEL Lowest-Observed-Adverse-Effect Level
31	miRNA MicroRNA
32	MEDINA Methylenedinitramine
33	MNX Hexahydro-1-nitroso-3,5-dinitro-1,3,5-triazine
34	MOA Mode of Action
35	NAS National Academy of Sciences
36	NCI National Cancer Institute
37	NIOSH National Institute for Occupational Safety and Health
38	NOAEL No-Observed-Adverse-Effect Level
39	NRC National Research Council
40	NTP National Toxicology Program
41	OECD Organization for Economic Cooperation and Development
42	ORD Office of Research and Development
43	OSF Oral Slope Factor
44	PBPK Physiologically Based Pharmacokinetic
45	PND Postnatal Day
46	POD Point of Departure

1	PTX	Picrotoxin
2	PWG	Pathology Working Group
3	RDX	Hexahydro-1,3,5-trinitro-1,3,5-triazine
4	RfC	Reference Concentration
5	ROS	Reactive Oxygen Species
6	RR	Relative Risk
7	SAB	Science Advisory Board
8	SDMS	Spontaneous Death or Moribund Sacrifice
9	TNX	Hexahydro-1,3,5-trinitroso-1,3,5-triazine
10	UCL	Upper Confidence Limit
11	UF	Uncertainty Factor
12	UF <sub>D</sub>	Database uncertainty factor
13	UF <sub>H</sub>	Human Inter-individual Variability Uncertainty Factor
14	UF <sub>L</sub>	LOAEL-to-NOAEL Uncertainty Factor
15	UF <sub>S</sub>	Subchronic-to-chronic Uncertainty Factor
16	WHO	World Health Organization
17		
18		

## 1. EXECUTIVE SUMMARY

The Science Advisory Board (SAB) was asked by the EPA's Integrated Risk Information System (IRIS) program to review the agency's *Draft IRIS Toxicological Review of Hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) (September 2016)* (hereafter referred to as the draft assessment). EPA's IRIS is a program that evaluates information on human health effects that may result from exposure to environmental contaminants. The draft assessment consists of a review of the available toxicologic scientific literature on RDX. The draft assessment was revised in September 2016 and a summary of EPA's disposition of the public comments received on an earlier draft version of the assessment was added to the Toxicological Review in Appendix E of the Supplemental Information.

### Literature Search Strategy/Study Selection and Evaluation

In general, the literature search strategy, study selection considerations, and study evaluation considerations, including inclusion and exclusion criteria, are well-described, documented, and appropriate. However, the SAB identified several areas that EPA's literature search missed and that should have been covered, including literature on the role of GABAergic systems in brain development and the potential developmental neurotoxicity of RDX through interference with GABAergic systems. In addition, in the literature search strategy section, EPA should clarify its reasoning and approach for including or excluding nonmammalian species studies and secondary references. The SAB notes that the metabolism of RDX has not been adequately studied, and suggests that the lack of toxicological data for the anaerobic bacteria metabolite, methylenedinitramine (MEDINA) and mammalian oxidative transformation product 4-nitro-2,4-diazabutanal (NDAB), and 4-nitro-2,4-diazabutanamide should be noted in the assessment. The SAB has identified additional peer-reviewed studies from the literature, which the agency should consider in the draft assessment.

### Toxicokinetic Modeling

The SAB finds the conclusions reached by the EPA following its evaluation of the PBPK models of Krishnan et al. (2009) and Sweeney et al. (2012a, b) to be well-documented and scientifically supported. The modifications that the agency made to the PBPK models of Krishnan/Sweeney represent distinct improvements over the original approach, and these changes adequately represent RDX toxicokinetics. The EPA also performed validation of the PBPK model using independent rat data sets, and all models provided reasonable fits according to standard goodness-of-fit measures. The SAB finds the uncertainties in the model to have been well described.

The SAB concludes that the choice of dose metric for neurotoxicity is clearly described. Without brain RDX concentration data, plasma or blood concentration data is used as a surrogate for brain concentrations. The agency's approach is adequately justified, since limited pharmacokinetic data in mice, rats, and swine and in humans show concordance between blood and brain RDX levels over time following exposure. The use of area under the curve (AUC) in a plasma concentration-time plot as a dose metric for interspecies extrapolation to humans from oral points of departure (PODs) derived from rat data is justified. AUC is representative of the

average RDX plasma concentration over a dosing interval, i.e., 24-hour interval. Published blood and brain RDX levels in rats for 24-hour time-courses appear to coincide with symptomatology. The mouse model was not used to derive PODs for noncancer or cancer endpoints because of uncertainties in the model as well as uncertainties associated with selection of a dose metric for cancer endpoints. This decision is scientifically supported and clearly explained.

## **Hazard Identification and Dose-Response Assessment**

### *Nervous System Effects*

The available human, animal, and mechanistic studies support EPA's conclusions that neurotoxicity, including seizures or convulsions, are human hazards of RDX exposure. Furthermore, RDX-induced convulsions arise primarily through a rapid mode of action resulting from RDX-induced blockade of the GABA<sub>A</sub> receptor (GABA<sub>A</sub>R) (Williams et al. 2011). Despite the limitations of the only cross-sectional study of Ma and Li (1993), which indicated significant neurobehavioral and memory deficits associated with RDX exposure for workers in a Chinese RDX plant, there is sufficient evidence from clinical case reports and experimental animals and mechanistic studies of RDX to support EPA's conclusion of neurotoxicity. Therefore, RDX should be considered for classification as a *potential convulsant to humans*. However, the SAB concludes that the evidence presented in the draft assessment does not adequately depict RDX's hazards to the nervous system. Convulsions in rodents can only provide a limited spectrum of potential human hazard; furthermore, convulsive or nonconvulsive seizures, epileptiform discharges, reduction in seizure threshold, subchronic sensitization, and neuronal damage can all be part of the spectrum of RDX's nervous system hazards. Additional future studies addressing cognitive and behavioral effects of RDX would assist in assessing other endpoints less severe than convulsions. Although there are data from existing animal studies showing changes in behavior, the data are not sufficiently robust to evaluate dose-response relationships, and animal data on cognitive changes is lacking. Therefore, there are no endpoints among existing studies to address the complete spectrum of RDX effects.

The SAB finds the selection of studies reporting nervous system effects to be scientifically supported and clearly described. It is appropriate to consider the dose-response data reported in Crouse et al. (2006) as a relevant model. While this study utilized gavage administration of RDX rather than a dietary route of administration, there is less variability in the amount of the toxic agent delivered by gavage compared to dietary intake and its dependence on the animal's feeding patterns. As long as these characteristics of administration are understood and accounted for, there is no reason to exclude work using gavage routes of exposure. The SAB agrees that the characterization of convulsions as a severe endpoint, and the potential relationship to mortality, is appropriately described.

The SAB finds that the selection of convulsions as the endpoint to represent nervous system hazard for RDX is scientifically supported and clearly described. Convulsion is the most biologically significant endpoint that has been reasonably and reliably measured for RDX. However, evidence from other seizurogenic compounds with a mode of action similar to RDX suggests that there are other, generally sub-clinical, cognitive and behavioral neurological effects that occur at doses below those causing seizure activity. The SAB agrees that the likely dose range between convulsion and other nervous system effects can be addressed using UF

adjustments. The SAB also finds that the calculation of the HEDs using PBPK modeling for the convulsion studies in rats to be scientifically supported and clearly described, and endorses the approach of estimating the effective concentration as the area under the curve (AUC) of concentration and time.

However, the SAB identifies several concerns regarding EPA's use of a BMR of 1% for benchmark dose modeling of the Crouse et al. (2006) data for convulsions. EPA's choice of a BMR of 1% for modeling is based on the severity of the convulsion endpoint and on the proximity (dose-wise) of convulsions to lethality. In the Crouse study, a BMR of 1% would correspond to a response that is a factor of 15 below the lowest observed response data. The SAB agrees that both the severity of convulsions as an endpoint and the proximity of convulsive doses to lethal doses are indeed valid sources of uncertainty in terms of providing sufficient protection for sensitive human populations. However, the SAB concludes that uncertainty about the appropriateness of the dose-response data and the POD derived from those data should be addressed through UFs and not through unsupported extrapolation of the dose-response data. A BMR of 5% based on the Crouse study is more consistent with the observed response at the Lowest-Observed-Adverse-Effect-Level (LOAEL) of 15% and not so far below the observable data. EPA should consider use of a 5% BMR with additional uncertainty factor to address the concern over using convulsions as the toxicological endpoint for the RfD. At a minimum, EPA will need to provide a more thorough justification for its choice of a 1% BMR, and specifically justify why a 1% BMR is a more appropriate extrapolation than a 5% BMR, and why the greater conservatism in risk assessment required for a frank effect is better dealt with through a lower BMR than through application of UFs.

With respect to the application of UFs to the PODs, the SAB supports the application of an interspecies UF of 3 to account for the toxicodynamic and residual toxicokinetic uncertainty in extrapolation from animal to human not accounted for by the toxicokinetic modeling, a LOAEL to No-Observed-Adverse-Effect (NOAEL) UF of 1, and an UF of 10 for intra-human variability. However, the SAB has concerns about the use of a subchronic to chronic UF (UF<sub>S</sub>) of 1. Observations in an *in vitro* assay for GABA activity found that the effects of RDX were not reversible following compound wash out (Williams et al. 2011). As such, there is the possibility that repeated exposures to RDX may have cumulative effects on GABAergic neurotransmission. The SAB recommends that EPA reconsider the UF for subchronic to chronic extrapolation, and at a minimum provide stronger justification for the use of a UF<sub>S</sub> of 1. In addition, the SAB questions the application of a database uncertainty factor (UF<sub>D</sub>) of 3, and recommends EPA consider applying a UF<sub>D</sub> of 10 to account for data gaps in developmental neurotoxicity, lack of incidence data for less severe effects, and proximity of the dose inducing convulsions to that inducing mortality. A composite UF of 300 should be considered instead of 100 as proposed in the draft assessment.

The SAB does not find the reference dose (RfD) derived by EPA for nervous system effects to be scientifically supported and concludes that the POD based on convulsions did not capture all of the potential adverse outcomes, or their severity. While the SAB recommends the assessment use the dose-response data of the Crouse et al. (2006) study as the primary basis for the derivation of an RfD for neurotoxicity, EPA should more fully account for database uncertainty.



1 *Kidney and other Urogenital System Effects*

2 The SAB agrees the available human, animal, and mechanistic studies support the conclusion  
3 that kidney and other urogenital system toxicities are a potential human hazard of RDX  
4 exposure. However, this conclusion is primarily supported by animal data, whereas available  
5 human studies identifying the kidney as a potential target of RDX are sparse and only identify  
6 transient renal effects following acute human exposure. There are no reports of prostatic effects  
7 of RDX in humans and no pertinent mechanistic data regarding RDX effects on the kidney and  
8 urogenital system. The SAB finds all hazards to the kidney and urogenital system adequately  
9 assessed and described in the draft assessment, with the exception of the description of  
10 inflammatory changes in the rat prostate. The SAB also finds that the selection of suppurative  
11 prostatitis as the endpoint to represent this hazard was clearly described in the draft assessment,  
12 but not scientifically supported because there is no known mechanistic link between suppurative  
13 prostatitis and renal papillary necrosis or adverse effects in the kidney.

14  
15 The SAB finds that the selection of the Levine et al. (1983) study on kidney and other urogenital  
16 system effects was clearly described, but not entirely supported by scientific evidence. Mild  
17 toxic effects of RDX exposure on the kidney were found in some species, but not others. In some  
18 studies, toxic effects were found in both sexes, while in others only male or female effects were  
19 observed. Of note is that some of these effects (i.e., mineralization) occurred in a small study  
20 with non-human primates, while some rodent studies did not find evidence of renal toxicity.  
21 Only in the chronic study of Levine et al. (1983) were severe toxic effects on the kidney found,  
22 and this was only seen in males at the highest dose (40 mg/kg-day); bladder toxicity also  
23 occurred in this treatment group, whereas effects on the prostate occurred at doses of 1.5 mg/kg-  
24 day and above. Therefore, the SAB determines that the selection of suppurative inflammation of  
25 the prostate as a “surrogate marker” of the observed renal and urogenital system effects for  
26 derivation of a reference dose to be not justified. The SAB recommends that a separate RfD be  
27 derived for renal papillary necrosis and the associated renal inflammation for the kidney and  
28 urogenital system and that the male accessory sex glands be designated as a separate organ  
29 system, with a separate RfD derived for suppurative prostatitis.

30  
31 As for calculation of the POD and HED for suppurative prostatitis as a stand-alone endpoint,  
32 both are scientifically supported and clearly described. The application of UFs should be the  
33 same as those for nervous system effects, if this system-specific RfD is to be considered for  
34 selection as an overall RfD.

35  
36 *Developmental and Reproductive System Effects*

37 The SAB disagrees with the conclusion in the draft assessment that there is suggestive evidence  
38 of male reproductive effects associated with RDX exposure. The available animal evidence  
39 based on testicular degeneration in male mice exposed to RDX in their diet for 24 months (Lish  
40 et al. 1984) is weak, unsupported by other endpoints in that study, complicated by the age of the  
41 mice and general toxicity of the RDX dose used, and contradicted by most other studies. In short,  
42 the database as a whole does not support this conclusion. There is no human evidence indicating  
43 male reproductive toxicity; no human studies have focused on this question, and there were no  
44 incidental reports of reproductive effects following RDX exposures. The SAB also finds  
45 adequate evidence from animal studies to conclude that RDX does not pose a risk of induction of  
46 structural malformations during human fetal development based on studies on rats and rabbits at

doses that were high enough to occasionally produce maternal toxicity. Additionally, the SAB agrees that conclusions cannot be drawn regarding other forms of developmental toxicity, which only occurred at maternally toxic dose levels. The SAB also concludes that RDX presents a potential neurodevelopmental hazard that was not adequately addressed in the draft assessment. A pilot developmental neurotoxicity study in rats found a significant concentration of RDX in the immature brain of offspring and in milk from dams treated with 6 mg/kg-day of RDX during gestation. Given that Lish et al. (1984) was used for the calculation of a POD and HED for the derivation of an organ/system-specific reference dose for reproductive system effects, the RfD based on testicular degeneration is not scientifically supported.

#### Other Noncancer Hazards

The SAB considers it important that the draft assessment be explicit as to whether the available evidence does or does not support liver, ocular, musculoskeletal, cardiovascular, immune, or gastrointestinal effects as a potential human hazard, and the rationale for reaching that conclusion. In addition, body weight gain should be included in this evaluation as it has been identified as a potential adverse effect of RDX exposure elsewhere.

#### Cancer

The SAB concurs with the EPA that “*suggestive evidence of carcinogenic potential*” is the most appropriate cancer hazard descriptor for RDX and that this descriptor applies to all routes of human exposure. The SAB agrees with the EPA that the relevant observations are the liver tumors that were observed in female B6C3F1 mice and male F344 rats and lung tumors that were observed in female B6C3F1 mice in two-year dietary bioassays (Lish et al. 1984; Levine et al. 1983). The SAB identifies a number of limitations for these studies and concludes that the evidence for a positive tumor response to RDX in two species, two sexes, or two sites, required by EPA’s *Guidelines for Carcinogen Risk Assessment* (USEPA, 2005) for a “*likely to be carcinogenic to humans*” descriptor, is weak or absent. On these bases, the SAB concludes that the descriptor, “*suggestive evidence of carcinogenic potential*”, is appropriate. The SAB also finds that the draft assessment adequately explains the rationale for a quantitative cancer dose-response analysis for RDX. Lish et al. (1984) was a well-conducted two-year bioassay that included a large number of animals tested at multiple dose levels, and increased incidences of neoplasms occurred in exposed female mice. The study is suitable and appropriate for dose-response assessment, consistent with EPA’s 2005 *Guidelines for Carcinogen Risk Assessment*.

With regard to the cancer dose-response assessment, the SAB supports the use of a linear low-dose extrapolation approach, as the mode of action for cancer resulting from RDX exposure is unknown. The SAB finds that the calculations of the PODs and oral slope factor are not clearly described, and the SAB questions whether these are scientifically supported. The SAB also has concerns with the unexpectedly low 1.5% incidence of liver tumors in female control mice and its impact on dose-response modeling. In addition, the draft assessment relies on the Multistage model to describe the POD and cancer slope factor. While understanding the preference of the IRIS program for the multistage model form, the SAB recommends that at a minimum, the draft assessment discuss the adequacy of the fit of the multistage model to the available data. The SAB also recommends that a more detailed description of the agency’s MS-COMBO modeling methodology be provided in the draft assessment to include a description of the independence assumption and the impact of violations of this assumption on the estimated POD.

## **Dose-Response Analysis**

### Oral Reference Dose for Effects Other Than Cancer

Although the SAB agrees that neurotoxicity should be the basis for an overall RfD for RDX, the SAB finds the scientific support for the proposed overall RfD to be weak. The SAB recommends that EPA use the dose-response data of the Crouse et al. (2006) study as the primary basis for the derivation of the overall RfD. An RfD derived from the NOAEL of Cholakakis et al. (1980) can be used for comparison.

### Inhalation Reference Concentration for Effects other than Cancer

There are no toxicokinetic (TK) data from inhalation exposures of laboratory animals or humans to RDX. There are epidemiological studies of persons exposed occupationally to RDX, but no information was provided on exposure levels. In light of the lack of TK data and exposure levels, an inhalation reference concentration cannot be derived.

### Oral Slope Factor for Cancer

The SAB finds that the calculation of an oral slope factor for cancer endpoints is not clearly described in the draft assessment, and has questions about whether the oral slope factor is scientifically supported. The SAB makes multiple suggestions on how the discussion can be improved.

### Inhalation Unit Risk for Cancer

There are no toxicokinetic data from inhalation studies of RDX in laboratory animals or humans, no inhalation carcinogenicity bioassays of RDX, nor data on cancer incidence in humans. Therefore, an inhalation unit risk for cancer cannot be derived.

## **Executive Summary**

Generally, the SAB considered the Executive Summary to be well written, succinct, and clear. As changes are made to the body of the draft assessment in response to the SAB's recommendations, the Executive Summary should be updated accordingly. In addition, the SAB offers a number of specific suggestions for improving the Executive Summary.

## 2. INTRODUCTION

The Science Advisory Board (SAB) was asked by the EPA Integrated Risk Information System (IRIS) program to review the agency's *Draft IRIS Toxicological Review of Hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX)* (hereafter referred to as the draft assessment). EPA's IRIS is a human health assessment program that evaluates information on health effects that may result from exposure to environmental contaminants. The draft assessment consists of a review of available scientific literature on RDX. The draft assessment was revised in September 2016 and a summary of EPA's disposition of the public comments received on an earlier version of the assessment was added in Appendix E of the Supplemental Information to the Toxicological Review.

In response to the EPA's request, the SAB convened an expert panel consisting of members of the Chemical Assessment Advisory Committee augmented with subject matter experts to conduct the review. The SAB panel held a teleconference on November 17, 2016, to discuss EPA's charge questions (see Appendix A), and a face-to-face meeting on December 12 - 14, 2016, to discuss responses to charge questions and consider public comments. The SAB panel also held teleconferences to discuss their draft reports on April 13, 2017, and April 17, 2017. Oral and written public comments have been considered throughout the advisory process.

This report is organized to follow the order of the charge questions. The full charge to the SAB is provided as Appendix A. Editorial comments from the SAB are provided in Appendix B. The SAB also provides suggestions on the format of EPA's charge questions in Appendix C.

### 3. RESPONSES TO EPA'S CHARGE QUESTIONS

#### 3.1. Literature Search/Study Selection and Evaluation

*Charge Question 1. The section on Literature Search Strategy/ Study Selection and Evaluation describes the process for identifying and selecting pertinent studies. Please comment on whether the literature search strategy, study selection considerations including exclusion criteria, and study evaluation considerations, are appropriate and clearly described. Please identify additional peer-reviewed studies that the assessment should consider.*

The literature search strategy, study selection considerations, and study evaluation considerations, including inclusion and exclusion criteria, are mostly well-described, documented, and appropriate, with a few exceptions noted below. EPA suitably cast a wide net to retrieve all pertinent studies for the evaluation of health effects associated with RDX exposure. They searched PubMed, Toxline, Toxcenter, Toxic Substances Control Act Test Submissions (TSCATS), and the Defense Technical Information Center (DTIC) database, a central online repository of defense-related scientific and technical information within the Department of Defense. Studies were then screened to find those relevant to assessing the adverse health effects of exposure to RDX and developing a dose-response assessment. Citations in review articles and citations within original articles were also obtained and screened for additional pertinent information.

Figure LS-1 and Table LS-1 provide a summary of the general inclusion and exclusion criteria for studies that were considered for further evaluation of potential health effects of RDX. EPA used criteria to exclude studies such as citations that were abstract only, on treatment and mitigation of environmental contamination with RDX, on laboratory methods, and those on the physical-chemical properties including explosivity. These were appropriate exclusion criteria, in the SAB's opinion. These criteria resulted in the exclusion of over 900 references from further evaluation. The SAB thought that Figure LS-1 could be clearer and better coordinated with the inclusion and exclusion criteria described in Table LS-1.

Table LS-1 indicates that studies on "ecological species" and nonmammalian species were also excluded. This contradicts statements (page xxix, lines 13-16) indicating that studies on nonmammalian species and ecosystem effects were considered as sources of information for the health effects assessment. The SAB suggests that these statements be clarified, and that data for all mammalian species be retained, even if they are considered "ecological species."

The SAB notes that the exclusion of nonmammalian species may not be appropriate in light of the use of nonmammalian species such as zebrafish (e.g., in medium throughput assays for developmental neurotoxicity) to evaluate potential health risk to humans, and describe Adverse Outcome Pathways. Although there may be no studies of RDX *in vitro* or in the cellular and tissue-based high throughput assays, future research using these types of assays may provide mechanistic information for chemicals that could be used in health effects assessments.

Inclusion criteria in Table LS-1 were related to whether a citation was a source of health effects data pertinent to assessing the risk to humans (e.g., studies of health outcomes in RDX exposed humans or standard mammalian models by either the oral or inhalation route; exposure to RDX measured; health outcomes/endpoints reported). Sources of mechanistic and toxicokinetic data

were also included. Secondary references and other sources that described ecosystem effects, exposure levels, dealt with mixtures, were reviews or risk assessments and regulatory documents, were excluded from study evaluation. However, EPA indicates that secondary references containing health effects data, and citations on nonmammalian toxicity were kept for consideration in the draft assessment. The description of what was done with secondary references could be clearer and better coordinated between the text and Figure LS-1 and Table LS-1.

EPA provides details of the search in Appendix B, including search terms, and the number of hits per search term sequence per database searched. They also tabulate the number of citations added to the database from their forward and backward web of science search of specific citations. Thus, the Agency has been transparent in its process of identifying studies for evaluation.

EPA's evaluation of studies is reasonably well-described and summarized in Table LS-3. The Agency used standard criteria and questions to evaluate study quality and utility that are described in several EPA guidance documents cited in the draft assessment. Studies were evaluated considering the experimental design and conduct, issues related to exposure to RDX, endpoints evaluated, and presentation of results. EPA describes generally the issues they considered in evaluating the utility of both human and animal studies to inform both hazard identification and dose-response assessment.

EPA excluded four studies on health effects and described the reason for excluding these in Table LS-2. Similarly, EPA describes some of the important limitations in experimental animal studies in Table LS-5. Overall, the description of EPA's study evaluation is clear, although the terminology is somewhat inconsistent (e.g., methodological features in Table LS-3 do not quite match the subheadings where these are described later in the section). Some details on strengths and limitations of specific studies chosen for further evaluation are provided in subsequent sections describing hazard identification and dose-response assessment for specific organ systems.

The SAB raised concerns about an inadequate description and discussion of supporting evidence for sensitive subpopulations in the draft assessment. Although there are no adequate studies on developmental neurotoxicity of RDX, there are some mechanistic studies implicating GABA antagonist activity of RDX in the neurotoxicity observed in animals and humans. The SAB concludes it would have been appropriate to search the literature for the role of GABA in brain development to describe what is known to date and incorporate this information into the draft assessment (see additional discussion of this issue in Section 3.3.1.4). Such mechanistic information provides evidence for the existence of sensitive subpopulations (e.g., infants, children, pregnant women and their fetus), and informs the choice of UFs meant to account for variability in the human population. EPA does not discuss the role of GABAergic systems in neurodevelopment and the potential for interference with this system by RDX (or other compounds with similar molecular mechanisms) to induce developmental neurotoxicity, an omission that should be rectified. The SAB identified six references that may be used to start the discussion of the role of GABAergic systems during development and the potential for RDX developmental neurotoxicity. A listing of these references is provided below.

The SAB notes that the metabolism of RDX has not been adequately studied. Limited toxicity information for the N-nitroso metabolites of RDX, specifically hexahydro-1-nitroso-3,5-dinitro-1,3,5-triazine (MNX) and hexahydro-1,3,5-trinitroso-1,3,5-triazine (TNX), has been discussed in the draft assessment and in the Supplemental Information document. However, reference to the anaerobic bacteria metabolite, methylenedinitramine (MEDINA) (Fuller et al. 2009 and 2010), has not been included in the metabolism section of the Supplemental Information document. The SAB suggests that the toxicity of MEDINA and other mammalian oxidative transformation products, e.g. 4-nitro-2,4-diazabutanal (NDAB), and 4-nitro-2,4-diazabutanamide be investigated in the future, as well as MNX and TNX be further investigated. N-nitroso metabolites are generated anaerobically and likely result from bacterial transformation of parent RDX in the gastrointestinal tract (Pan et al. 2007b). Although these are minor metabolites, some reductive transformation products of RDX (including MNX and TNX) are present in ground waters near munitions and training facilities (Beller and Tiemeier, 2002),

The SAB assembled seven additional references to augment the neurotoxicity database. In addition, 11 candidate references that address the production and toxicity of reductive transformation products and studies that were conducted in species that may inform the current RDX assessment are identified. A full listing of these references is provided below.

***Key Recommendations:***

- EPA should include a literature search on the role of GABAergic systems in brain development, and how this knowledge can inform a better understanding of the potential developmental neurotoxicity of RDX.
- EPA should not exclude nonmammalian species as they may bring important mechanistic insight into the draft assessment.
- EPA should clarify its reasoning and approach for including or excluding nonmammalian species studies and secondary references.

***Suggested Recommendations:***

- The lack of / paucity of toxicological data for MEDINA and the mammalian oxidative transformation product 4-nitro-2,4-diazabutanal (NDAB), 4-nitro-2,4-diazabutanamide, MNX and TNX should be noted in the draft assessment.

***Future Needs:***

- More research on the metabolism of RDX to identify metabolites and their potential toxicity is needed.
- Adequate developmental neurotoxicity studies are needed given the mechanism of action of RDX.

***Additional Citations for USEPA to Consider:***

1. Beller, HR; Tiemeier, K. (2002). Use of liquid chromatography/tandem mass spectrometry to detect distinctive indicators of in situ RDX transformation in contaminated groundwater. *Environmental Science & Technology* 36: 2060-2066.

2. Creeley, CE. (2016) From drug-induced developmental neural apoptosis to pediatric anesthetic neurotoxicity – where are we now? *Brain Sci* 6(3):32-44.
3. Fuller, ME; Perreault, N; Hawari, J. (2010). Microaerophilic degradation of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) by three *Rhodococcus* strains. *Letters in Applied Microbiology* 51:313–318.
4. Fuller, ME; McClay, K; Hawari, J; Paquet, L; Malone, TE; Fox, BG; Steffan, RJ. (2009). Transformation of RDX and other energetic compounds by xenobiotic reductases XenA and XenB. *Appl Microbiol Biotechnol* 84:535-544.
5. Gust, KA; Brasfield, SM; Stanley, JK; Wilbanks, MS; Chappell, P; Perkins, EJ; Lotufo, GR; Lance, RF. (2011). Genomic investigation of year-long and multigenerational exposures of fathead minnow to the munition compound RDX. *Environ Toxicol Chem* 30: 1852-1864.
6. Halasz, A; Manno, D; Perreault, NN; Sabbadin, F; Bruce, NC; Hawari, J. (2012). Biodegradation of RDX Nitroso Products MNX and TNX by Cytochrome P450 XplA. *Environ Sci Technol* 46: 7245-7251.
7. Jaligama, S; Kale VM, Wilbanks, MS, Perkins, EJ, Meyer, SA. (2013). Delayed myelosuppression with acute exposure to hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) and environmental degradation product hexahydro-1-nitroso-3,5-dinitro-1,3,5-triazine (MNX) in rats. *Toxicol Appl Pharmacol* 266: 443-451.
8. Jeilani, YA; Duncan, KA; Newallo, DS; Thompson, AN, Jr.; Bose, NK. (2015). Tandem mass spectrometry and density functional theory of RDX fragmentation pathways: Role of ion-molecule complexes in loss of NO<sub>3</sub> and lack of molecular ion peak. *Rapid Commun Mass Spect* 29: 802-810.
9. Kim, JY; Liu, CY; Zhang, F; Duan, X; Wen, Z; Song, J; Feighery, E; Lu, B; Rujescu, D; St Clair, D; Christian, K; Callicot, JH; Weinberger, DR; Song, H; Ming, GL. (2012). Interplay between DISC1 and GABA signaling regulates neurogenesis in mice and risk for schizophrenia. *Cell* 148:1051-1064.
10. Marty, S; Wehrle, R; Sotelo, C. (2000). Neuronal activity and brain-derived neurotrophic factor regulate the density of inhibitory synapses in organotypic slice cultures of postnatal hippocampus. *The Journal of Neuroscience* 20:8087-8095.
11. Meyer, SA; Marchand, AJ; Hight, JL; Roberts, GH; Escalon, LB; Inouye, LS; MacMillan, DK. (2005). Up-and-down procedure (UDP) determinations of acute oral toxicity of nitroso degradation products of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX). *J Appl Toxicol* 25: 427-434.
12. Mukhi, S; Patino, R. (2008). Effects of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) in zebrafish: General and reproductive toxicity. *Chemosphere* 72: 726-732.



13. Neal, AP; Guilarte, TR. (2013). Mechanism of lead and manganese neurotoxicity. *Toxicol Res* 2(2):99-114.
14. Rivera, C; Voipio, J; Payne, JA; Ruusuvuori, E; Lahtinen, H; Lamsa, K; Pirvola, U; Saarma, M; Kaila, K. (1999). The K<sup>+</sup>/Cl<sup>-</sup> co-transporter KCC2 renders GABA hyperpolarizing during neuronal maturation. *Nature* 397(6716):251-5.
15. Salari, AA, Amani, M. (2017) Neonatal blockade of GABA-A receptors alters behavioral and physiological phenotypes in adult mice. *Int J Dev Neurosci* 57:62-71.
16. Smith, JN; Pan, XP; Gentles, A; Smith, EE; Cox, SB; Cobb, GE. (2006). Reproductive effects of hexahydro-1,3,5-trinitroso-1,3,5-triazine in deer mice (*Peromyscus maniculatus*) during a controlled exposure study. *Environ Toxicol Chem* 25: 446-451.
17. Williams, LR; Wong, K; Stewart, A; Suciu, C; Gaikwad, S; Wu, N; DiLeo, J; Grossman, L; Cachat, J; Hart, P; Kalueff, AV. (2012). Behavioral and physiological effects of RDX on adult zebrafish. *Comparative Biochemistry and Physiology C-Toxicology and Pharmacology* 155:33-38.
18. Wirbisky, SE; Weber, GJ; Lee, JW; Cannon, JR; Freeman, JL. (2014) Novel dose-dependent alterations in excitatory GABA during embryonic development associated with lead (Pb) neurotoxicity. *Toxicology Letters* 229:1-8.

### **3.2. Toxicokinetic Modeling**

In Appendix C, Section C.1.5, the draft assessment presents a summary, evaluation, and further development of published PBPK models for RDX in rats, mice, and humans (Sweeney et al. 2012a; Sweeney et al. 2012b).

#### **3.2.1. Model Evaluation**

*Charge Question 2a. Are the conclusions reached based on EPA's evaluation of the models scientifically supported? Do the revised PBPK models adequately represent RDX toxicokinetics? Are the model assumptions and parameters clearly presented and scientifically supported? Are the uncertainties in the model appropriately considered and discussed?*

The conclusions reached by the EPA following its evaluation of the PBPK models of Krishnan et al. (2009) and Sweeney et al. (2012a, b) are well-documented and scientifically supported. EPA did a thorough and accurate job reviewing and summarizing much of what is known about the oral absorption of different forms/preparations of RDX, as well as the compound's distribution, metabolism and excretion. The changes that the agency made to the PBPK model of Krishnan /Sweeney represent distinct improvements over the original approach, and these changes adequately represent RDX toxicokinetics. Human metabolic rate constants were fitted from human data. Additionally, it is stated that *in vitro* data from rats and human metabolic studies were used and scaled-up to liver size based on microsomal protein. The EPA also performed validation of the PBPK model using independent rat data sets, and the models provided reasonable fits according to standard goodness-of-fit measures. The uncertainties in the model

are well-described and were appropriately considered as illustrated by the discussion of the mouse model and the decision not to implement it. Overall, the SAB finds that the model assumptions and parameters were scientifically supported and that the draft assessment does an excellent job in compiling the data presented in Appendix C.

The SAB has several suggestions based on its review of Section C.1:

- In Section C.1.2, include the tissue parent and metabolite data of Pan et al. (2013) cited elsewhere in the report.
- In Section C.1.2., provide more text describing the distribution of RDX to the brain as a key target tissue. Issues that could be discussed in more detail include i) Brain extracellular fluid concentration-effect relationships; ii) Changes in plasma/blood concentrations over time may be linearly related (proportional) to brain concentrations, and may be used to derive toxicity, as proposed, based on limited correlations observed with brain and plasma data from animal studies and data from a child after poisoning (Woody et al. 1986); iii) what led to the decision to not use PBPK-simulated brain RDX concentrations, which were only moderately well fitted in Figure C-6, as a dosimeter for neurotoxicity risk assessment; and iv) Experimental findings that lend support to the decision to use plasma as a surrogate.
- Protein binding of RDX is not mentioned in the draft assessment. This may be regarded as a potential weakness given that it is the free concentration that would diffuse across the blood-brain barrier in the absence of any active uptake processes, or be available for metabolism. This could lead to differences in predicted brain/blood ratios in humans, and may be helpful in allometric scale-up of clearance. However, absent any empirical values for protein binding, use of total, rather than free, concentrations is the only option.
- Despite improvements in the model, the rat data are only moderately well fitted and show substantial deviations, especially at early time-points. This may reflect deviations of the simulations due to inaccurate model absorption parameters, and possibly imprecise clearance parameters. Further optimization may improve fitting. Insight into the nature of gastrointestinal absorption could be gained from *in vitro* studies using Caco2 cells or other intestinal models. For elimination, hepatic intrinsic clearance is preferred over a rate constant. From the *in vitro* microsomal and S9 studies reported by Cao et al. (2008), data are provided that can be used to calculate metabolic intrinsic clearance. The Cao study demonstrated that the intrinsic metabolic clearance in a microsomal preparation was greater in humans than in rats and mice. However, concentration-dependent studies were not performed, so this publication does not provide support for the assumption of linear clearance.
- Clearance terms instead of first order rate constants (dependent on elimination and the apparent volume of distribution) would be more informative in the model. *In vitro* ( $K_m/V_{max}$  or intrinsic metabolic clearance) or derivation of intrinsic clearance from fitted clearance obtained from *in vivo* data may be used.
- The role of metabolites in toxicity is discussed in the draft assessment, but due to a lack of data not included in the model. This is appropriate, though limited information on metabolites in brain and other tissues (Pan et al. 2013) indicates they could contribute to the observed effects. The parent AUC dose metric would thus serve as an indicator of exposures to parent and metabolites, though not directly tracking the metabolites.

- Provision for tissue partitioning is mainly via *in silico* methods; more *in vivo* data would provide justification for these values.
- The mouse data are the least comprehensive, though EPA could re-evaluate whether the total radioactivity data in Guo et al. (1985) are consistent with the Sweeney et al. (2012) data. The description of the noncompartmental analysis (page C-28) is good to provide perspective on the data versus the PBPK modeling.

**Key Recommendations:**

- The current modeling as performed is reasonable and appropriate, based on the currently available pharmacokinetic data.

**Suggested Recommendations:**

- Revise the text to address the issues listed above as warranted.

**3.2.2. Selection of Dose Metric**

*Charge Question 2b. The average concentration of RDX in arterial blood (expressed as area under the curve) was selected over peak concentration as the dose metric for interspecies extrapolation for oral points of departure (PODs) derived from rat data. Is the choice of dose metric for each hazard sufficiently explained and appropriate? The mouse PBPK model was not used to derive PODs for noncancer or cancer endpoints because of uncertainties in the model and because of uncertainties associated with selection of a dose metric for cancer endpoints. Is this decision scientifically supported?*

For neurotoxicity, the choice of dose metric is clearly described (pages 2-8 and 2-9); however, the basis for the choice of dose metric for the prostatitis endpoint should be described. The choice is reasonable and appropriate, given less than ideal data on the pharmacokinetic-pharmacodynamic (PK/PD) relationship for this endpoint. A PK/PD model likely would be driven by the concentration in brain that is responsible for the PD (neurotoxicity); brain RDX concentrations are derived from the blood-brain partitioning of RDX blood concentrations. Without brain RDX concentration data, plasma or blood is used as a surrogate for brain concentrations. The agency's approach is adequately justified and appropriate, since limited PK data in mice, rats, and swine (Table C-1) and in a human (Woody et al. 1986) show concordance between blood and brain RDX levels over time following exposure, supporting the use of blood/plasma concentrations as a surrogate for brain concentrations, and for the use of plasma concentration-time curve AUC values as a dose metric.

AUC is representative of the average RDX plasma concentration over a dosing interval, i.e., 24-hour interval. Published 24-hour time courses of blood and brain RDX levels in rats (e.g., Bannon et al. 2009) appear to coincide with symptomatology, providing support for the use of AUC. It is appropriate to assume that seizures or hyperreactivity would be manifest as long as a threshold blood/brain concentration of RDX, e.g., 8 µg/g (Williams et al. 2011) is present. Therefore, there is clear rationale for choosing AUC over peak plasma concentrations ( $C_{max}$ ) values as the dose metric.

The POD<sub>HED</sub> is presented in Table 2-2 of the draft assessment for both dose metrics for neurotoxicity, with the difference between C<sub>max</sub> and AUC/24 hour values being relatively modest in the rat (~30%). It should also be pointed out in the text on pages 2-8 that AUC appears to be a better representation of the adverse effect of interest than is RDX concentration at a single point in time. Additionally, it should be noted that maximal plasma concentrations are not predicted well from the PBPK model, producing uncertainty in C<sub>max</sub> values, and supporting the case for the use of AUC.

There does not appear to be an explanation for the choice of dose metric for the prostatitis endpoint, though some comments (e.g., AUC considered better estimated than C<sub>max</sub> from PBPK model) in the discussion for neurotoxicity apply across endpoints. Again the differences in Table 2-2 are modest, and since this is an effect only observed in a chronic study, average daily AUC is an appropriate choice.

It is noted that although there are mechanistic data to support the role of RDX in neurotoxicity (convulsions) through binding to GABA<sub>A</sub>R (Williams et al. 2011; Williams and Bannon, 2009), the effect of RDX might be due to either parent compound or metabolites; as such, any PK parameter that measures parent compound plasma concentrations may not accurately predict toxicity.

The mouse PBPK model was not used to derive PODs for noncancer or cancer endpoints because of uncertainties in the model and because of uncertainties associated with selection of a dose metric for cancer endpoints. This decision is scientifically supported and clearly explained on pages C-30 and C-31. The mouse model is more uncertain as discussed on pages 2-9 of the draft assessment. The uncertainties are numerous.

**Key Recommendation:**

- Current approaches for dose metrics are appropriate, but the basis for the choice of dose metric for the prostatitis endpoint should be described.

**3.2.3. Intrahuman Variation**

*Charge Question 2c. In Section 2.1.3 of the draft assessment, an uncertainty factor of 10 for human variation is applied in the derivation of the RfD. Does the toxicokinetic modeling support the use of a different factor instead?*

It is standard practice to adopt an intraspecies factor of 10 to account for potential differences in the toxicokinetics and toxicodynamics of a chemical in the absence of information about variability within human populations. There is a paucity of data on the toxicokinetics, toxicodynamics or toxicity of RDX in humans. Given these extreme data limitations and the likely toxicodynamic and toxicokinetic differences, it would not be appropriate to use a PBPK model and it is appropriate to use a full UF<sub>H</sub> of 10.

Sensitivity analyses (described in Appendix C) showed that the PBPK model output was substantially impacted by bioavailability and by metabolic clearance. There are apparently no data to define the absorption phase following RDX ingestion by humans or animals. Toxicokinetic data for RDX elimination by humans are quite sparse. It appears from two studies

(Bhushan et al. 2003; Major et al. 2007) that RDX metabolism in some mammals is mediated by cytochrome P450s (CYPs). As the activities of CYPs and other enzymes that metabolize xenobiotics vary significantly in the human population, the rate of metabolic clearance of RDX would also be expected to vary. Potential inter-subject differences in formation of RDX metabolites may also contribute to uncertainty, should specific metabolites be associated with toxicities.

In light of the role of binding of RDX to the GABA<sub>A</sub>R in neurotoxicity, data on inter-subject variability in receptor binding and response could identify and characterize sensitive subpopulations.

**Key Recommendation:**

- The likely toxicodynamic and toxicokinetic differences support a full UF<sub>H</sub> of 10.

**Future Needs:**

- Data on inter-subject variability in receptor binding and response are needed.

**3.3. Hazard Identification and Dose-Response Assessment**

**3.3.1. Nervous System Effects**

**3.3.1.1. Nervous System Hazard**

*Charge Question 3a(i). The draft assessment concludes that nervous system toxicity is a human hazard of RDX exposure. Please comment on whether the available human, animal, and mechanistic studies support this conclusion. Are all hazards to the nervous system adequately assessed? Is there an appropriate endpoint to address the spectrum of effects?*

The SAB agrees that available human, animal, and mechanistic studies support the conclusion that nervous system toxicity is a human hazard of RDX exposure.

**Human Studies**

There is consistent evidence from more than 20 clinical case reports that exposure to RDX is associated with adverse neurological outcomes, particularly convulsions. Nevertheless, there are many and varied limitations to deducing the hazards of RDX solely based on such case reports. There is only one cross-sectional study that provides a snapshot of the potential neurotoxicity associated with inhalation exposure to RDX. Ma and Li (1993) presented results from a neurobehavioral test battery that also assessed memory retention in workers exposed in a Chinese RDX plant at a single point in time. The results indicated that significant neurobehavioral and memory deficits were associated with RDX exposure measured in air. However, this study has several significant limitations that impact any conclusions about RDX hazard solely based on its findings. SAB's greatest concerns with this study are: 1) the omission of exposure levels in the "non-exposed" group; 2) no attempt to control for confounders (non-occupational exposures, lifestyle, co-morbidity), and; 3) no rationale provided for subdividing the exposed cohort. Nevertheless, the outcomes on Composite Memory Retention Quotient and Composite Block Design score were greater than 15 points and greater than 2 seconds lower

( $p < 0.01$ ) than the control group, respectively. The statistical analyses performed seems appropriate, but 95% confidence intervals would have been helpful given that the magnitude of functional impairments across groups is within the High Average [110-119] and Average [90-109] range, measures typically associated with a 15% Standard Deviation. Other studies are generally supportive, with the strongest evidence for convulsions coming from investigations involving acute exposures (Testud, 1996; Hollander, 1969; Merrill, 1968).

#### Animal Studies

Several studies with rodents using oral gavage and dietary exposure over the acute (Burdette 1988), sub-chronic (Crouse et al. 2006; Von Oettingen, 1949) and chronic (Lish 1984, Levine 1983, Hart 1976) timeframes have consistently identified a broad range of neurological impairments, ranging in severity from irritability to tremors and other signs that may be considered prodromal of convulsions. Convulsive (seizure) activity is a common finding in most, but not all, studies. In addition to seizure activity, several of these studies (Levine et al. 1990; Angerhofer et al. 1986; Levine et al. 1983; Levine et al. 1981; von Oettingen et al. 1949) identified “less severe” neurological and behavioral impairments (e.g. hyperactivity and nervousness) that may be consistent with findings identified in the sparser literature on human exposures. Some of the animal study findings suggest that RDX appears to sensitize animals exposed at lower doses to subsequent seizurogenic stimuli, including electrogenic, audiogenic, and chemical kindling.

#### Mechanistic Studies

The neurotoxicity profile of RDX is consistent with that of a centrally acting excitotoxicant. There is ample evidence of a direct interaction of RDX with GABA<sub>A</sub>R in the mammalian central nervous system. RDX blocks GABA-activated chloride ion currents and the inhibitory postsynaptic potentials (IPSPs) that form critical inhibitory networks throughout the brain. The available data do not preclude the influence of other unstudied receptors for RDX, but also implicate the limbic system, including the amygdala, as particularly sensitive targets of RDX. Surprisingly, the potency of RDX as a GABA<sub>A</sub>R blocker is relatively low. By comparison, picrotoxin (PTX) has a greater than 100-fold lower  $K_i$  (100x more potent) than RDX at binding to GABA<sub>A</sub>R with a  $K_i$  (inhibition constant) of 0.2 vs. 21  $\mu$ M (Williams et al. 2011). The lower potency of RDX extends to the concentrations needed to inhibit chloride ion currents in whole cell voltage clamp experiments and inhibitory postsynaptic current (IPSC) events, which typically require greater than 10  $\mu$ M. Also relevant to the RDX mechanism and its potential importance to long-term behavioral toxicity is the observation that the inhibitory actions of RDX on seizure-like neuronal discharges can be measured in the basolateral nucleus of the amygdala (Williams et al. 2011). In contrast, evidence supporting a direct role for glutamate in the effects of RDX is limited, and a basis for excessive glutamate stimulation in the draft assessment is weak, if not unfounded. However, the overall excitation within neuronal networks of the adult mammalian brain is controlled primarily, though not exclusively, by the balance of glutamatergic (excitatory) and GABAergic neurotransmission among interconnected circuits, and thus are inextricably linked.

#### Conclusions

Regarding nervous system hazard identification, the available human, animal, and mechanistic studies support EPA’s conclusions that neurotoxicity, including seizures or convulsions, are

1 human hazards of RDX exposure. Furthermore, RDX-induced convulsions arise primarily  
2 through a rapid mode of action resulting from RDX-induced GABA<sub>A</sub>R blockade. Despite the  
3 limitations of the Ma and Li (1993) study, the sum of the evidence from clinical case reports,  
4 experimental animals, and mechanistic studies of RDX indicates there is sufficient evidence to  
5 support this conclusion. Therefore, RDX should be considered for classification as a *potential*  
6 *convulsant to humans* who are at risk for exposures to RDX.

7  
8 The evidence presented in the draft assessment, however, does not fully depict RDX's hazards to  
9 the nervous system. The SAB notes that convulsions in rodents only provide a limited spectrum  
10 of potential human hazard, with convulsive or non-convulsive seizures, epileptiform discharges,  
11 reduction in seizure threshold, subchronic sensitization, and neuronal damage all being part of  
12 the spectrum of RDX's nervous system hazards. Further evaluation or explanation should be  
13 provided in the report for these additional potential endpoints. With respect to whether all  
14 hazards to the nervous system were adequately assessed, the measure of abnormal electrographic  
15 activity or seizure-like activity in specific brain regions may be a more sensitive indicator of  
16 neurotoxicity than the potential of RDX to elicit subtler neurological impairments such as  
17 cognitive deficits and/or behavioral abnormalities.

18  
19 Although endpoints such as convulsions, tremors and aggression are appropriate as part of the  
20 spectrum of effects, it is important to note that the functional observation battery (FOB) data  
21 presented in Crouse et al. (2006) are rudimentary and not sufficiently sensitive to detect  
22 neurobehavioral consequences produced by chronic/subchronic doses of RDX over prolonged  
23 periods, especially during pregnancy. Moreover, tests designed to detect subtle developmental  
24 neurotoxicity during the perinatal-weaning period have not been conducted in any studies to  
25 date. Most of the FOB methods used were observational and highly descriptive, and are  
26 considered blunt instruments likely to have missed relevant neurological impairments, if present.  
27 The data presented by Crouse et al. (2006) sets a range and NOAEL for convulsion, but no  
28 conclusions can be reached about the lower limit of sub-convulsive doses that are without frank  
29 neurological impacts, including fine psychomotor impairments, anxiety and social impairments,  
30 decreased executive functioning and long term memory. These concerns are compelling because  
31 of more recent peer-reviewed published data indicating that sub-convulsive doses of either  
32 bicuculline (which has a similar mechanism of action to RDX) or domoic acid (which has  
33 agonist activity on glutamate transmission) have been shown to cause developmental and  
34 behavioral impairments at doses below those that cause convulsions (Salari and Amani, 2017;  
35 Gill et al. 2010). GABAergic and glutamatergic neurotransmission are inextricably linked, not  
36 only in regulating excitability of the adult brain, but their balance throughout perinatal  
37 development provides essential developmental cues that refine functional neural connectivity.  
38 The lack of scientific information about the influences of RDX on this balance is a major  
39 uncertainty. Thus, the SAB concludes that there remains significant uncertainty about the  
40 developmental neurotoxicity of RDX. Additional studies addressing cognitive and behavioral  
41 effects of RDX would assist in assessing other endpoints less severe than convulsions. Although  
42 there are data from existing animal studies showing changes in behavior, the data are not  
43 sufficiently robust to evaluate dose-response relationships, and animal data on cognitive changes  
44 are lacking. Given these limitations, additional studies measuring other neurological endpoints  
45 are needed to address the complete spectrum of effects.

**Future Needs:**

- Provide a means to commission behavioral and morphometric studies that can permit more accurate assessments of RDX exposures to the developing nervous system at subconvulsive doses.
- Conduct studies with adequate power to address cognitive and behavioral effects, as well as developmental neurotoxicity of RDX, and to establish relevant dose-response relationships.

**3.3.1.2. Nervous System-Specific Toxicity Values**

*Charge Question 3a(ii). Please comment on whether the selection of studies reporting nervous system effects is scientifically supported and clearly described. Considering the difference in toxicokinetics between gavage and dietary administration (described in Appendix C, Section C.1, and in the context of specific hazards in the toxicological review), is it appropriate to consider the Crouse et al. (2006) study, which used gavage administration? Is the characterization of convulsions as a severe endpoint, and the potential relationship to mortality, appropriately described?*

The selection of studies reporting nervous system effects is scientifically supported, clearly described, and provides sufficient information to identify the central nervous system as a primary toxicological target for exposures to RDX. Based on a review of the scope of the search strategy and the process for identifying studies that report health effects and meet appropriate standards of quality for conduct, design, and reporting, the SAB concludes that the most reliable scientific information has been accessed for this draft assessment.

For assessment of nervous system effects, the convulsion endpoint is appropriate for revealing the hazards of RDX delivered by oral gavage administration (Crouse et al. 2006; Cholakakis et al. 1980) or the dietary route (Levine et al. 1983; Lish et al. 1984).

However, the available data from animal studies with RDX do not adequately address the potential effect of low-level exposure(s) of RDX either through adult life stages, or most importantly, during the highly susceptible perinatal period (see Section 3.3.1.1). Based on the current state of the science with other better studied compounds known to elicit seizures by similar or functionally related mechanisms, the SAB cannot discount whether RDX is capable of producing subtle, yet relevant, behavioral, psychomotor, or cognitive outcomes. The SAB generally agrees that the balance/imbalance of neuronal network excitation/inhibition during the perinatal and post-weaning periods of development have profound and measurable influences on the functional and anatomical integrity of developing neuronal networks. This not only impacts behavioral, psychomotor, and cognitive outcomes throughout the lifespan, but can also promote significantly greater susceptibility to subsequent exposures to other seizurogenic chemicals or physical triggers of convulsion (Stamou et al. 2013; Lee et al. 2016; Meunier et al. 2017). There is a wealth of peer-reviewed experimental evidence showing that even modest impairments in the excitation/inhibition balance of developing neuronal circuits, whether originating from genetic mutations, chemical exposures, or their combination can effect long-lived (possibly permanent) changes in behavioral, psychomotor, and cognitive outcomes. This is particularly



important for chemicals that interfere with GABAergic neurotransmission. It is also important to emphasize the developmental transition of GABA<sub>A</sub> receptor from excitatory to inhibitory neurotransmission during the perinatal/postnatal period, which adds additional complexity to how dose and timing of RDX exposures alters neurological outcomes. This is especially important since RDX has been shown to interact in a competitive manner with picrotoxin at GABA<sub>A</sub> receptors of basolateral amygdala (BLA), but once it alters their function, its actions are not reversible (Williams et al. 2011).

Therefore, there are major gaps in our knowledge about exposures to sub-convulsive doses of RDX and their possible neurological ramifications, especially during the perinatal and early weaning periods of development. Clearly, such exposures are possible and relevant, and could have consequences not only to individuals directly exposed to RDX, but also those exposed transplacentally and/or during lactation. Additional developmental neurotoxicity studies need to be conducted in animals to address these gaps, including test batteries to detect potential fine psychomotor impairments, anxiety and social impairments, decreased executive functioning and long-term memory.

Biological plausibility for such detrimental actions comes from a rich literature demonstrating that developmental exposure *in vivo* and *in vitro* to seizurogenic chemicals, albeit through distinct molecular mechanisms, have potent influences on outcomes relevant to developmental neurotoxicity at concentrations below those that elicit convulsions, but influence behavioral, psychomotor, and cognitive outcomes. Examples relevant to RDX include the GABA<sub>A</sub>R antagonist bicuculline (Grasso et al. 2016; Nasehi et al. 2017; Salari et al. 2017), and domoic acid (Costa et al. 2010; Doucette et al. 2016; Grant et al. 2010; Hiolski et al. 2016; Marriott et al. 2016; Mills et al. 2016; Zuloaga et al. 2016). Finally, Zhang and Pan (2009) provided strong evidence that adult mice fed diets with RDX at a sub-convulsive dose of 5 mg/kg for 28 days resulted in significant changes in key miRNA brain transcripts that are related to neurological and metabolic functions, and also were changed in a tissue-specific manner. It is appropriate to conclude that such effects are likely to exert long-lived consequences on brain development should they be elicited in the perinatal period (Hu and Li, 2017).

The differences in toxicokinetics of RDX exposure by gavage versus dietary administration are clear, and must be accounted for when predicting risk. Animal studies reporting effects on neurological health utilized gavage or dietary route of administration. The evidence indicates that the gavage route results in higher peak blood and brain levels of RDX than the dietary route and that the rate of rise in blood and brain levels is faster with gavage. Gavage results in more reliable and consistent dose to blood and brain than dietary intake. Moreover, incidences of convulsions were not reported in most dietary studies (Levine et al. 1983; von Oettingen et al. 1949). The SAB concludes that it is appropriate to consider the dose-response data reported in the Crouse et al. (2006) study as a relevant model. In fact, the Crouse study produced perhaps the best RDX dose-response data available for convulsion.

EPA also used the incidence data for convulsions in pregnant dams dosed by gavage with RDX from gestation days 6 – 19 from the teratology study of Cholakakis et al. (1980) for dose-response assessment. The candidate POD and RfD derived from the Cholakakis et al. study was 5 times lower than those derived using data from the subchronic study of Crouse et al. (2006). This may

1 indicate that pregnancy is a sensitive window for neurotoxicity in the adult. Or, it may indicate  
2 that the higher sample size of 24 to 25 per dose used by Cholakakis et al. (1980) was sufficient to  
3 detect convulsions at a lower dose than the Crouse study, which had a sample size of 10 animals  
4 per dose. However, EPA noted limitations in the Cholakakis et al. study concerning quantification  
5 of the dose-response relationship and elected to use the Crouse study as the basis of the proposed  
6 RfD value. The SAB supports this decision, as detailed below.

7  
8 The noted limitations in the Cholakakis study included a lower purity test compound, a 14- day  
9 dosing regimen rather than 90 days, and three widely spaced (order of magnitude) dose  
10 groupings versus five tightly spaced dose groupings in the Crouse study, all of which impact the  
11 accuracy of a POD. Although the test substance purity was lower in the Cholakakis study, it is not  
12 clear how much difference this would make in the study results given that after subtracting out  
13 water, the purity in Cholakakis was 90% compared to 99.9 % in the Crouse study. Another  
14 limitation cited for the Cholakakis et al. (1980) study was difficulty keeping the dosing suspension  
15 uniformly suspended. In both the Cholakakis et al. (1980) teratology study and the Crouse study,  
16 doses of RDX were administered in a methyl cellulose / Tween 80 vehicle as a suspension.  
17 Assay results for the dosing suspensions presented in the appendix of the Cholakakis report  
18 demonstrate high variability and the study authors acknowledged “maintaining uniform  
19 suspensions was not always easy.” When the same nominal concentration was assayed  
20 repeatedly, it showed wide variation in RDX content (33% to 500% relative to nominal),  
21 although the RDX concentration of one of the assayed dosing suspensions was much higher  
22 (500%) than nominal and skewed the range of variability. Most assayed dosing suspensions  
23 were lower than nominal. Less variability in RDX dose suspensions was observed in the Crouse  
24 study, in which each dose suspension was mixed using a magnetic stirring bar until a uniform  
25 suspension was obtained, and continued to be mixed each day during the dosing procedure. Since  
26 these measures were taken by Crouse et al. to reduce the variation in dosing suspensions, it is  
27 likely that the intended dose levels were more accurately administered in the Crouse study  
28 compared with the Cholakakis study. The SAB recognizes that the study limitations increase  
29 uncertainty in deriving the RfD based on the Cholakakis et al. (1980) study. However, the SAB  
30 also notes that Cholakakis et al. (1980) observed convulsions in a pregnant dam at a dose (2 mg/kg-  
31 day) lower than the LOAEL in the Crouse study (8 mg/kg-day) in male and nonpregnant female  
32 rats. Further, in the Angerhofer et al. (1986) teratology study, one death was reported in the  
33 dams at 2 mg/kg-day and one death at 6 mg/kg-day, although the authors did not report whether  
34 convulsive symptoms occurred prior to death. Nonetheless, there is some suggestion, although  
35 the data are weak, that pregnancy may be a sensitive physiological state for the neurotoxicity of  
36 RDX. Overall, considering all of the above factors, the SAB concludes that it is appropriate to  
37 give more weight to the Crouse study with respect to the quantitative dose-response analysis.

38  
39 The SAB agrees that the characterization of convulsions as a severe endpoint, and its potential  
40 relationship to mortality, are appropriately described. Based on the available data, death may  
41 occur without seizure or convulsions, although this may simply be due to a low frequency of  
42 observations. However, based on the current state of science (including the epilepsy literature),  
43 death is not a necessary outcome of seizures or convulsions, and is driven by abnormal  
44 electrographic patterns in the brain. While the relationship between convulsions and mortality is  
45 unclear in the overall scheme of assessment of neurotoxicity endpoints for RDX, it is nonetheless  
46 appropriate to conclude that convulsions, as characterized in the draft assessment, represent a

reasonable severe endpoint for human health risk assessment. In addition, more consideration should be given to available data on fatal outcomes and the possibility that mortality may arise from non-nervous system factors or hazards.

***Suggested Recommendation:***

- More consideration should be given to discussing available data on fatal outcomes and the possibility that mortality may arise from non-nervous system factors or hazards.

***Future Needs:***

- The SAB strongly recommends developmental neurotoxicity studies to be conducted in animals. These studies should include test batteries to detect potential fine psychomotor impairments, anxiety and social impairments, decreased executive functioning and long-term memory.

**3.3.1.3. Points of Departure for Nervous System Endpoints.**

*Charge Question 3a(iii). Is the selection of convulsions as the endpoint to represent this hazard scientifically supported and clearly described? Are the calculations of PODs for these studies scientifically supported and clearly described? Is the calculation of the HEDs for these studies scientifically supported and clearly described? Does the severity of convulsions warrant the use of a benchmark response level of 1% extra risk? Is calculation of the lower bound on the benchmark dose (BMDL) for convulsions appropriate and consistent with the EPA's Benchmark Dose Guidance?*

**Convulsions Endpoint:**

The SAB finds that the selection of convulsions as the endpoint to represent nervous system hazard for RDX is scientifically supported and clearly described. The evidence indicates that convulsions are the most biologically significant endpoint that has been reasonably and reliably measured. However, the SAB notes that evidence from other seizurogenic compounds with similar modes of action suggests that there are other, generally sub-clinical cognitive and behavioral neurological effects that occur at lower doses. It is likely that such effects also occur for RDX, although data to establish this are not currently available. For compounds such as bicuculline, triggering of abnormal biochemical, electrographic patterns and/or abnormal connectivity measured by magnetic resonance imaging (MRI) or positron emission tomography (PET) approaches can be demonstrated to occur at doses below those that cause seizures (Bruyns-Haylett et al. 2017; Galineau et al. 2017; Nasrallah et al. 2017; Takahashi et al. 2017). As such, the SAB agrees that the likely dose range between convulsion and other nervous system effects can be addressed using the UF adjustments.

**POD Calculations:**

The draft assessment determined that the incidence data for convulsions from Crouse et al. (2006) and Cholakakis et al. (1980) were amenable to BMD modeling. PODs based on a BMR of 1% extra risk for convulsions were calculated for both studies. In addition, a POD based on the NOAEL for convulsion from the two-year dietary study of Levine et al. (1983) was also derived. The SAB concludes that the PODs for convulsions were clearly described and correctly calculated. However, the SAB questions the use of BMR of 1% extra risk for convulsions. In

1 addition, the SAB's detailed discussion on the appropriateness of the Crouse study for the RfD  
2 and hence estimation of the POD is provided in its response to charge question 4a. presented in  
3 Section 3.4.1. The SAB's discussion on the choice of a BMR of 1% is provided later in this  
4 section.

#### 5 6 HED Calculations:

7 The SAB agrees that the calculation of the HEDs for these studies is scientifically supported and  
8 clearly described. EPA estimates the HED by assuming the equivalent pharmacokinetic potency  
9 of equivalent rat and human arterial blood concentrations of RDX. The concentration of RDX as  
10 a function of time following dosing is generated using a PBPK model, and the effective  
11 concentration is estimated as the AUC of concentration and time. The SAB endorses this  
12 approach. The SAB agrees that, given the binding of the parent compound to the GABA<sub>A</sub>R, a  
13 dose metric for the parent compound is appropriate, though it also may be serving as a surrogate  
14 if any metabolites also have that activity. The AUC is a more appropriate choice than C<sub>max</sub> to  
15 estimate the effective concentration due to the uncertainties in the parameterization of the model  
16 for absorption.

#### 17 18 Benchmark Response Level for Convulsions:

19 The SAB identifies the following concerns regarding EPA's use of a BMR of 1% in the  
20 benchmark dose model of the dose-response data from Crouse et al. (2006). Based on EPA's  
21 *Benchmark Dose Technical Guidance* (U.S.EPA, 2012a), the "standard reporting level"  
22 (although not *per se* the default) BMR for quantal data (such as those for the incidence of  
23 convulsions) is 10%. In that guidance, EPA suggests conditions that would justify BMR values  
24 less than 10%. The justification given in the guidance for applying a smaller BMR is "biological  
25 considerations" of the endpoint being modeled. EPA's guidance does, in fact, identify "frank  
26 effects" as an example of such biological considerations for choosing a BMR of "5% or lower."  
27 In addition, a 1% BMR is recommended for epidemiological data. However, the guidance also  
28 points out that "...if one models below the observable range, one needs to be mindful that the  
29 degree of uncertainty in the estimates increases. In such cases, the BMD and BMDL can be  
30 compared for excessive divergence. In addition, model uncertainty increases below the range of  
31 data." In its clarification of this choice to the SAB, the EPA stated that the BMR of 1% was  
32 chosen based on biological considerations as given in its Benchmark Dose Technical Guidance.  
33 Specifically, EPA stated that this BMR was chosen to address the fact that the endpoint being  
34 modeled, in this case convulsions is a frank effect. The SAB acknowledges that the convulsions  
35 observed by Crouse et al. (as well as by Cholakakis et al.) is indeed, a frank effect, and the SAB is  
36 sensitive to the need to provide an adequate margin of safety to protect against even a low  
37 frequency of occurrence of this effect. However, the SAB finds that EPA's Benchmark Dose  
38 Technical Guidance is vague on how "biological considerations" should influence the  
39 benchmark dose modeling. The lack of clarity in the use of this term, and the absence of  
40 guidance as to how "biological considerations" should be applied in choosing a BMR, makes its  
41 application subjective.

42  
43 Benchmark dose modeling was developed to address the constraints placed upon dose-response  
44 assessment by the use of only study-specific, and dose-specific NOAELs and LOAELs, and to  
45 fully utilize the data on response to dosing. The purpose of benchmark dose modeling is to  
46 derive PODs from study data that are more generalizable to the inherent dose-response of a given

chemical than are the study's NOAEL or LOAEL. Benchmark dose modeling is viewed primarily as a process for modeling the dose-response *per se* using few, if any, assumptions that are extraneous to the data (except in rare cases where mechanistic information may inform the shape of the dose-response curve). Consistent with this view, the BMR should be strongly linked to the nature of the dose-response data. Hence, the caution expressed in EPA's benchmark dose guidance that, "...if one models below the observable range, one needs to be mindful that the degree of uncertainty in the estimates increases. In addition, model uncertainty increases below the range of data." Thus, the BMR should be close to (although not necessarily within) the observable data. The BMR determines the "distance" between the observable data and the BMD. As indicated in the EPA guidance document, the greater the "distance" between the observable data and the BMD, the greater the statistical uncertainty in the fit of the model at the BMD and, therefore, the greater the difference between the BMD and the BMDL. The computations presented in Table 1 show that the response at the LOAEL for the Crouse et al. (2006) study is 15% and for the Cholakakis et al. study is 4%. Thus, a BMR of 1% corresponds to a response that is a factor of 15 below the lowest observed responses for the study chosen for estimation of the POD.

**Table 1. LOAELs and Percent Response at LOAELs for Crouse et al. (2006) and Cholakakis et al. (1980)**

Study	n/dose group	LOAEL	Percent response at LOAEL
Crouse et al. (2006)	10 rats/sex/dose – group	8 mg/kg/d	15%
Cholakakis et al. (1980)	24-25 pregnant rats/dose group	2 mg/kg/d	4%

Table 2 below presents the BMDs for the Crouse et al. (2006) and Cholakakis et al. (1980) studies that would result from BMRs of 1%, 5% and 10%. The EPA benchmark dose guidance suggests looking at the BMD/BMDL ratios, also provided in Table 2, resulting from each BMR but provide no guidance on what constitutes a ratio indicative of unacceptable statistical uncertainty, and thus a BMR too low to be supported by the data. Note that the BMDL at a BMR of 1%, the EPA estimated POD, is roughly 4 times smaller than the BMDL at a BMR of 5%.

**Table 2. Comparison of BMDs and BMDLs at different BMRs for Crouse et al. (2006) and Cholakis et al. (1980)**

Study	BMR	BMD (mg/kg-d)	BMDL (mg/kg-d)	BMD/BMDL
Crouse et al. (2006) LOAEL = 8 mg/kg-d	1%	3.02	0.569	5.3
	5%	5.19	2.66	2.0
	10%	6.60	4.59	1.4
Cholakis et al. (1980) LOAEL = 2 mg/kg-d	1%	0.179	0.123	1.5
	5%	0.915	0.628	1.5
	10%	1.88	1.29	1.5

In its benchmark dose modeling of the data on the occurrence of convulsions in the Crouse et al., (2006) data, the EPA has attempted to address two different considerations, the dose-response relationship *per se*, and the frank nature of the effect. As discussed above, while EPA's own Benchmark Dose Technical Guidance provides for incorporating consideration of the frank nature of an effect into benchmark dose modeling, the details of how this best accomplished are unclear. However, the guidance cautions against overly diverging from the empirical data.

Therefore, EPA needs to revisit the choice of BMR and, at a minimum, provide a more thorough justification for its choice of a 1% BMR for the Crouse et al. data, including why a 1% BMR is an appropriate extrapolation below the available data and why a 1% BMR is a more appropriate extrapolation than use of a 5% BMR based on the Crouse et al. (2006) data for RDX.

Additionally, the SAB recommends that EPA provide a discussion and justification for why the greater conservatism in risk assessment required for a frank effect is better dealt with through a lower BMR than through application of UFs.

#### BMDL for Convulsions:

The calculation of the lower bound on the benchmark dose (BMDL) for convulsions is appropriate and consistent with EPA's Benchmark Dose Guidance. For the parameters specified by EPA (including the choice of a BMR of 1%), the benchmark dose is calculated per EPA's benchmark dose guidance. The choice of the model from among the available dose-response models is appropriate.

**Key Recommendations:**

- EPA should consider using a BMR of 5% for their dose-response modeling of the Crouse et al. (2006) data while addressing the uncertainty of using data on a frank effect (convulsions in this case) as the basis of an RfD with additional uncertainty factors.
- If EPA decides to use a BMR of 1% for the dose-response assessment using Crouse et al. (2006), EPA should justify why the greater conservatism in risk assessment required for a frank effect (due to the lack of incidence data for less severe endpoints) is better dealt with through a lower BMR than through application of UF<sub>D</sub>.
- If EPA decides to use a BMR of 1% for the Crouse et al. (2006), EPA should provide in its discussion clear justification for why a 1% BMR is more appropriate than a 5% BMR for RDX, given the greater uncertainty introduced into the dose-response assessment for RDX using a BMR of 1%.

**3.3.1.4. Uncertainty Factors for Nervous System Endpoints**

*Charge Question 3a(iv). Is the application of uncertainty factors to these PODs scientifically supported and clearly described? The subchronic and database uncertainty factors incorporate multiple considerations; please comment specifically on the scientific rationale for the application of a subchronic uncertainty factor of 1 and a database uncertainty factor of 3?*

EPA applied Benchmark Dose Software models to data from two gavage studies in rats (Crouse et al, 2006; and Cholakakis et al. 1980) to derive benchmark dose for a 1% response rate (BMDL<sub>01</sub>) as a point of departure for effects on the nervous system, following Human Equivalent Dose (HED) adjustment. A third data set (Levine et al. 1983) in rats was evaluated using the NOAEL approach. The toxicological endpoint in all cases was convulsions. EPA applied UFs to the HEDs to derive the proposed RfD for nervous system effects.

*Interspecies Uncertainty Factor (UF<sub>A</sub>)*

An interspecies uncertainty factor, UF<sub>A</sub>, of 3 ( $10^{1/2} = 3.16$ , rounded to 3) was applied to the points of departure (PODs), in this case the human equivalent dose for a 1% response rate, to account for the toxicodynamic and residual toxicokinetic uncertainty in extrapolating from average animal models to average humans not accounted for by the toxicokinetic modeling. This is standard risk assessment practice where an adequate toxicokinetic model was applied to derive a human equivalent dose, and available data are not sufficient to define quantitative toxicodynamic differences between species. The SAB agrees that the UF<sub>A</sub> of 3 is appropriate and clearly described.

*Subchronic to Chronic Uncertainty Factor (UF<sub>S</sub>)*

EPA chose the subchronic study of Crouse et al. (2006) to derive a RfD for nervous system effects and a UF<sub>S</sub> of 1 to extrapolate from a subchronic experimental exposure duration to chronic exposure. The SAB has concerns about the use of a UF<sub>S</sub> of 1. An *in vitro* assay of GABA activity showed that the effects of RDX were not reversible following compound wash out (Williams et al. 2011). Furthermore, in a 14-day range-finding study in Sprague Dawley rats, Crouse et al. (2006) observed convulsions (incidence and severity not reported) at doses of 17 mg/kg-day and above, but in the 90-day study on F344 rats, convulsions were elicited at 8 mg/kg-day (male: 1/10; female: 2/10) and above. Though the reported increase in sensitivity to

convulsions in the latter study may be due to the larger number of animals per group (10 animals/group vs. 6 animals/group in the 14-day study), the finding of convulsions at the lower dose in the 90-day study by the same investigators using the same procedures for administering RDX may reflect the influence of the longer exposure period. Thus, it is possible that chronic exposures to RDX may lead to progressive, possibly additive, inhibition of GABA<sub>A</sub>R in the brain. These observations raise the possibility of progressive, possibly cumulative effects on GABAergic neurotransmission that are not predicted by either RDX pharmacokinetics in the blood nor by the total levels of RDX in the brain, as only a minor fraction of total RDX in the brain would be bound to the GABA<sub>A</sub>R pool. There may be some compensation in the balance of excitatory and inhibitory neurotransmission, but the potential of such compensation to mitigate effects of RDX and the impact of compensation on the organism are unclear.

The SAB also has concerns about part of the EPA's rationale for using a UF<sub>s</sub> of 1, namely that "in studies of subchronic or gestational exposure used to derive a POD, effects were seen at lower doses in the studies of shorter duration than in the chronic studies" (Section 2.1.1, p 1-11 in the draft assessment). The three studies used to generate PODs were a gestational study with 14-day gavage exposure to pregnant dams (Cholakakis et al. 1980), a 2-year dietary study in male and female rats (Levine et al. 1983), and a 13-week gavage study in male and female rats (Crouse et al. 2006). As discussed in Section 3.3.1.2, pregnant dams in Cholakakis et al. gestational study may be a potential sensitive subpopulation, and are not readily comparable to non-pregnant animals. Thus, the 14-day Cholakakis study and 90-day Crouse study should not be compared to evaluate the effect of exposure duration on convulsant dose.

As EPA notes in the discussion of the studies and in Appendix C, differences in the method of dose administration, the physical form of RDX, including particle size, and/or dose matrix in the dietary studies and gavage preparations might influence the rate of absorption and internal dose, and may partly explain the differences in neurotoxic symptoms reported in the studies of varying duration, both dietary and gavage. RDX administered orally as a coarse particle preparation was shown to be more slowly absorbed than as a fine particle preparation (Schneider et al. 1977), thus influencing the kinetics of RDX. Differences in particle size in the 13-week dietary study in mice of Cholakakis et al. (RDX particle size of about 200 µm), and the 2-year dietary study in the same strain of mice by Lish et al. (RDX particle size less than 66 µm), may partly explain why no convulsions were reported at higher doses in the 13-week study, but convulsions were observed at lower doses in the 2-year study. However, differences in study population, dosing preparations, route of administration, and other methodological considerations make comparisons across studies for the purpose of evaluating effect of exposure duration difficult, if not impossible. Thus, EPA's statement that effects were seen at lower doses in shorter duration exposures is inappropriate. In making comparisons of the toxicity of RDX after different durations of exposure, these factors (e.g., particle size, dosing method, and dose matrix) that are known to influence rate of gastrointestinal absorption and/or bioavailability, should be addressed where possible. Note that the test material used in the key study of Crouse et al. (2006), although noted to be of higher purity than most other studies, was not characterized with respect to particle size.

The SAB recognizes that the NOAEL for convulsions in the 2-year dietary study in rats (Levine et al. 1983) was 8 mg/kg-day, which was 2-fold higher than the NOAEL of 4 mg/kg-day for



convulsions in the 13-week gavage study in rats (Crouse et al. 2006), and may be the basis for the application of a  $UF_s$  less than the default value of 10. However, the differences in the observed convulsant doses may be due to differences between dietary and gavage administration. As discussed in Section 3.3.1.2, RDX administered via gavage results in more reliable and consistent dose to blood and brain than dietary intake. Moreover, the 2-year dietary study was not designed for dose-response assessment for convulsions in exposed animals, and incidences of convulsions were not reported. Thus, occurrences of convulsions may have been missed during the course of the study.

Thus, the SAB recommends that EPA reconsider the UF for subchronic to chronic extrapolation, and at a minimum provide stronger justification for a  $UF_s$  of 1. Binding of RDX to the  $GABA_A$ R may provide for cumulative effects on inhibitory neurotransmission. Further, the uncertainty in doses received by animals in the various studies due to particle size and related issues makes cross-study comparison of the effects of duration of exposure inappropriate.

#### LOAEL to NOAEL Uncertainty Factor ( $UF_L$ )

The  $UF_L$  is meant to account for uncertainties in extrapolating from a LOAEL to a NOAEL when estimating an RfD. EPA applied a  $UF_L$  of 1 because the BMDL was used as a point of departure in Crouse et al. (2006) and in Cholakakis et al. (1980), and a NOAEL was used as the point of departure in Levine et al. (1983). Thus, no extrapolation from a LOAEL to a NOAEL was needed. This is standard risk assessment practice and the Panel agrees that this choice is appropriate and clearly described.

#### Database Uncertainty Factor ( $UF_D$ )

The EPA applied a  $UF_D$  of 3 in developing an RfD based on neurotoxicity to help account for database deficiencies. The SAB has several concerns regarding the large amount of database uncertainty for RDX including lack of developmental neurotoxicity testing, frank effect as a basis for the RfD with no available incidence data for less severe neurotoxicity, and proximity of the dose inducing convulsions with that inducing mortality.

The SAB is concerned that there is limited information available to understand developmental neurotoxicity of RDX. Transplacental and lactational transfer of RDX in rodents has been observed (Hess-Ruth et al. 2007), and therefore, there is potential exposure to the developing fetus and infant from maternal exposure. It is worth noting that Hess-Ruth et al. (2007) concluded that developmental neurotoxicity studies should be conducted for RDX, but apparently, this has not been done. EPA noted that the two-generation reproductive and developmental toxicity study of Cholakakis et al. (1980) did not report effects in the offspring at doses lower than maternally toxic doses. However, the study only looked at histopathology of the F2 pups at weaning. This study did not assess developmental neurotoxicity in the offspring. The draft assessment indicates that the existing literature did not demonstrate early life stage as a sensitive subpopulation, but this was not fully evaluated in animal studies and cannot be evaluated with the available human data. There was one case report involving one child poisoned by RDX, but this one case study does not provide evidence regarding the influence of age at exposure on toxicity.

RDX interferes with neurotransmission by binding at the GABA<sub>A</sub>R, and acting as an antagonist inhibiting GABAergic neurotransmission. GABA is a major inhibitory neurotransmitter in the adult brain. However, GABAergic systems play another role in vertebrate brain development acting as an excitatory neurotrophic factor contributing to processes involved in neurodevelopment (see for example Rivera et al. 1999; Kim et al. 2012). Lead (Pb), a potent developmental neurotoxicant, acts at least partly by inhibiting the GABAergic system during development in zebrafish (Wirbisky et al. 2014) and in mammals (reviewed in Neal and Guilarte, 2013). There is evidence that exposure of early postnatal rodent hippocampal slices to a GABA antagonist (bicuculline) reduces GABAergic neuroactivity, affects the regulation of GABAergic inhibitory synapses and increases their density in the hippocampus (Marty et al. 2000). The hippocampus is involved in seizure development in humans with epilepsy, so these results seem pertinent. There is evidence that drugs that act through the GABA<sub>A</sub>R as GABA agonists can also cause neurodevelopmental disorders (see for example the review by Creeley, 2016). These lines of evidence point to potential window(s) of susceptibility in the developing brain to chemicals interfering with GABAergic systems. Additional discussion is provided in Section 3.3.1.2.

As discussed in Section 3.3.1.2, there are major gaps in our knowledge about exposures to sub-convulsive doses of RDX and their possible neurological ramifications, especially during the perinatal and early weaning periods of development. Clearly, such exposures are possible and relevant, and could have consequences not only to individuals directly exposed to RDX but also those exposed transplacentally and/or during lactation.

Additional evidence prompting concern for developmental neurotoxicity is found in the section of the draft assessment on the mode of action of RDX neurotoxicity. The draft assessment cites a study (Zhang and Pan, 2009) reporting that RDX upregulates 3 microRNAs that affect brain-derived neurotrophic factor (BDNF) in the brains of mice fed 5 mg RDX/kg diet (estimated doses 0.75 to 1.5 mg/kg-day; Bannon et al, 2009). As EPA notes, BDNF is a member of the neurotrophin family of growth factors, and promotes the survival and differentiation of existing and new neurons. BDNF is important for brain development. Disrupting the regulation of BDNF could result in developmental deficiencies in the brain if the perturbation of this pathway were strong enough. This provides additional indirect evidence to generate concern for potential developmental neurotoxicity of RDX. Zhang and Pan (2009) provided strong evidence that adult mice fed RDX at a sub-convulsive dose of 5 mg/kg for 28 days resulted in significant tissue-specific changes in key miRNA brain transcripts related to neurological and metabolic functions. It is appropriate to conclude that such effects could have long-lived consequences on brain development should they be elicited in the perinatal period (Hu 2017). EPA does not discuss the role of GABAergic systems in neurodevelopment and the potential for interference with this system by RDX (or other compounds with similar molecular mechanisms) to induce developmental neurotoxicity, an omission that should be rectified (see recommendation under section 3.1). Until there are adequate developmental neurotoxicity studies on this compound, the potential for developmental neurotoxicity as an outcome of RDX exposure remains a significant data gap.

The SAB notes that EPA chose to model a BMDL<sub>01</sub> rather than a BMDL<sub>05</sub> because of the severity of convulsions as an endpoint. However, while recognizing the severity of the convulsion endpoint, the SAB, as discussed in its response to Charge Question 3a(iii) (Section

3.3.1.3), has concerns about the use of a BMR of 1% because of the degree of uncertainty it introduces in the benchmark dose analysis, and suggests that EPA considers the use of a BMR of 5%. While recognizing the need for conservatism in the development of the RfD given the use of a frank effect (convulsions) as the critical effect, the SAB suggests that rather than trying to capture this conservatism in the benchmark dose analysis, the EPA consider whether the  $UF_D$  is the more appropriate framework to provide this protection. Further, choosing a lower BMR from a study in adult animals does not account for the potential of widely different toxicodynamics by age at exposure.

EPA's BMD modeling (see Appendix D in Supplemental Information for the draft RDX assessment) of the mortality data also indicates that convulsive doses and lethal doses are approximately the same. The  $BMDL_{01S}$  for lethality from studies amenable to modeling overlay the  $BMDL_{01S}$  for convulsions. Note that the Crouse et al. (2006) study authors state that their study provides a NOAEL of 4 mg/kg-d for lethality. This is the same NOAEL for convulsions. Thus, mortality occurs in the same dose range as convulsions. This finding provides additional compelling support for using a  $UF_D$  of 10 rather than of 3.

Given the potential for neurodevelopmental toxicity of RDX through interference with GABAergic systems and other pathways, the proximity of lethal doses to convulsive doses, and the lack of incidence data on less severe neurotoxic effects of RDX, the SAB strongly recommends that EPA use a  $UF_D$  of 10 rather than 3. However, it should be noted that animals in both the Crouse et al. (2006) study and the Cholakis et al. (1980) gestational exposure study were exposed via gavage. The peak blood concentrations resulting from a bolus dose are expected to be higher than those from exposure via oral ingestion of contaminated water or food. Thus, there is some unquantified margin of safety in using a gavage study as the basis for an RfD that will be applied to typical human exposure scenarios.

#### Intraspecies Uncertainty Factor ( $UF_H$ )

EPA applied an intraspecies uncertainty factor of 10 to account for toxicokinetic and toxicodynamic variability in the human population. Although a PBPK model was used to extrapolate from the animal internal dose (AUC of RDX in arterial blood) to a human equivalent dose, EPA noted that not enough toxicokinetic data were available from human studies to quantify differences among humans.

The SAB agrees that the  $UF_H$  needs to account for both toxicodynamic and toxicokinetic variability among humans, and that not enough data were available to quantify toxicokinetic or toxicodynamic differences among humans. EPA used the standard default  $UF_H$  of 10, which is typically viewed as a composite of a half-log for toxicokinetic differences and a half-log for toxicodynamic differences. Toxicokinetic differences among humans can be related to age, pregnancy, illness, medication use, other chemical exposures, and so on. In the absence of adequate toxicokinetic data to model the range of differences among humans, such differences must be accounted for by including a default toxicokinetic component in the  $UF_H$ . The default toxicodynamic portion of the  $UF_H$  accounts for differences in target tissue or receptor-mediated response across humans. As noted above, there is limited evidence (Cholakis et al. 1980) that pregnant rats may be more sensitive to RDX than non-pregnant rats. The intraspecies UF is meant to account for differences across the human population. Pregnant animals represent one

potential sensitive subpopulation. For reasons stated above, the SAB agrees the use of a  $UF_H$  of 10 is scientifically supported and clearly described.

### **Key Recommendations**

- Consistent with EPA guidance for UFs, the SAB strongly suggests applying a  $UF_D$  of 10 to account for data gaps for developmental neurotoxicity, lack of incidence data for less severe neurological effects resulting in use of a severe effect (convulsions) as a basis for the RfD, and the proximity of lethal doses to convulsive doses.
- EPA should discuss whether potential neurodevelopmental effects of RDX would be sufficiently addressed by a  $UF_D$  of 10.
- SAB recommends that EPA reconsider the UF for subchronic to chronic extrapolation, and at a minimum provide stronger justification for a  $UF_S$  of 1.

### **Future Needs**

- Adequate studies on RDX to fully assess the risk of neurotoxicity as a developmental outcome of RDX exposure.
- Studies to determine whether pregnant rats are more sensitive to RDX than non-pregnant rats.
- More definitive study to look at effects of exposure duration that can better inform subchronic to chronic extrapolation.

#### **3.3.1.5.Nervous System-specific Reference Dose**

*Charge Question 3a(v). Is the organ/system- specific reference dose derived for nervous system effects scientifically supported and clearly characterized?*

Regarding the RfD for nervous system effects, the point-of-departure derived from the neurotoxicity assessment, based on convulsions as the critical endpoint, does not capture all potential adverse outcomes or their severity. This is one reason the SAB recommends increasing the  $UF_D$  to 10 (see Section 3.3.1.4). Recognizing the study quality concerns in Cholakakis et al. (1980), particularly with respect to the accuracy of administered doses, the draft assessment should utilize the dose-response data of Crouse et al. (2006) as the primary basis for the RfD. The POD from the observations in the Crouse study (see response to charge question 4a) is considered to be more reliable and accordingly should be given more weight.

Overall, the conclusion that the available data in humans and animals support a convulsant neurotoxicity for RDX, possibly through GABA<sub>A</sub>R blocking mode of action, is scientifically-supported. The proposed nervous system-specific reference dose was clearly described but not scientifically supported.

### **Key Recommendation**

- EPA should utilize the dose-response data of Crouse et al. (2006) as the primary basis for the RfD.
- The SAB recommends increasing the  $UF_D$  from 3 to 10.
- The SAB recommends revisiting the  $UF_S$  and providing a better justification, at a minimum, for the use of a  $UF_S$  of 1.

## **Future Needs**

- Additional dose specifications (levels) should be examined to provide a more reliable dose-response relationship for convulsion and other neurotoxic effects of RDX.

### **3.3.2. Kidney and Other Urogenital System Effects**

#### **3.3.2.1. Kidney and Other Urogenital System Hazard** (Sections 1.2.2, 1.3.1)

*Charge Question 3b(i). The draft assessment concludes that kidney and other urogenital system toxicity is a potential human hazard of RDX exposure. Please comment on whether the available human, animal, and mechanistic studies support this conclusion. Are all hazards to kidney and urogenital system adequately assessed? Is the selection of suppurative prostatitis as the endpoint to represent this hazard scientifically supported and clearly described?*

#### Available Human, Animal, and Mechanistic Studies:

The available human, animal, and mechanistic studies support the conclusion that toxicity to the kidney and other components of the urogenital system is a potential human hazard of RDX exposure. However, this conclusion is primarily supported by animal data, whereas available human studies that implicate the kidney as a potential target of RDX are sparse and only identify transient renal effects following acute human exposure. There are no reports of prostatic effects of RDX in humans and no pertinent mechanistic data regarding RDX effects on the kidney and urogenital system.

#### Hazards to Kidney and Urogenital System:

All hazards to the kidney and urogenital system are adequately assessed and described in the draft assessment, with the exception of the description of inflammatory changes in the rat prostate. The description in the draft assessment of these prostatic inflammatory changes should include not only suppurative inflammation, but also chronic inflammation and the variability and uncertainty in the classification of prostatic inflammation.

#### Selection of Suppurative Prostatitis Endpoint:

The selection of suppurative prostatitis as the endpoint (“surrogate marker”) to represent this hazard is clearly described in the draft assessment, but not scientifically supported because of various uncertainties that are associated with the hazard, including the following:

- There is no known biological or mechanistic basis for using suppurative prostatitis as a surrogate marker for renal and other urogenital (GU) effects.
- There is uncertainty about the direct association between suppurative prostatitis and the toxic renal effects observed in male rats in the Levine et al. (1983) study. A strong association between kidney lesions (papillary necrosis, pyelonephritis, and peri-renal peritonitis) and suppurative inflammation in the prostate was observed only in male rats in the highest dose group (40 mg/kg-day); there were no such renal changes in the lower dose groups (except in one male animal in the 8.0 mg/kg-day group), while suppurative prostatitis occurred at the two next highest doses (8.0 and 1.5 mg/kg-day). The renal lesions were considered primary effects of RDX in the draft assessment, while the prostatitis was considered secondary to the renal effects; the SAB concurs with this notion.

- There are uncertainties regarding the diagnosis of suppurative inflammation:

- (a) Suppurative and chronic inflammation are part of a continuum, and diagnostic criteria may have varied over time and among pathologists. Prostatic inflammation found in aged rats is divided into several subtypes, only one of which is suppurative inflammation. Other categories include subacute inflammation, chronic-active inflammation, and micro-abscesses. Reference is made in the draft assessment to a paper by Suwa et al. (2001) on the background pathology in the prostate of 1,768 control F344 rats allowed to live for up to 2.4 years. This paper was the basis for the conclusion by EPA that inflammation in the control group of the study by Levine et al. (1983) was unusually low for this strain of rat. However, in the paper by Suwa et al. (2001), all types of inflammation are combined, and 70.4% of these rats had inflammation mostly confined to the dorsolateral prostate and graded as mild. No data were provided by Suwa et al. on suppurative inflammation.
- (b) Combining all types of prostate inflammation in the 24-month groups of the Levine et al. (1983) study yields similar incidences among all groups, with the exception of the highest dose group. The prostatitis incidences in the control and the three lowest dose groups were about 40% lower than the incidences reported by Suwa et al. (2001) for aged F344 rats in NTP studies; the lower incidences may be a reflection of the manner of histopathologic examination (see point c below).

In the 24-month study by Levine *et al.* (1983), 22/54 control rats (41%) had prostatic inflammation of any type (chronic and suppurative inflammation) and the incidence in the 0.3, 1.5, and 8.0 mg/kg-day dose groups was 22/55 (40%), 22/52 (42%), and 23/55 (42%), respectively; and these lesions were mostly graded as minimal to mild. Two of the three rats that died between six and 12 months in the 1.5 mg/kg-day group had subacute prostatic inflammation, and subacute prostatitis was also diagnosed in one of 10 rats in the 1.5 mg/kg-day dose group and two of 10 rats of the 8.0 mg/kg-day dose group that were euthanized at the scheduled 12-month interim time point.

By contrast, 51 of 55 rats in the high dose (40 mg/kg-day) group of the 24-month study died before the end of the two-year study and 39 of these 51 rats (76%) that died had prostatic inflammation. Prostatic inflammation (chronic type) was found in one of five rats in the high dose group that died before six months and in 18 of 19 (95%) rats in the high dose group that died between six and 12 months. Suppurative inflammation was found in 19 of 27 (70%) rats that died between 12 and 24 months in the high dose group and chronic inflammation occurred in one of these 27 rats. One of the four high dose rats that lived until the end of the study had minimal chronic inflammation of the prostate; no suppurative prostatic lesions were found in these four rats. Thus, the total incidence of prostatic inflammation of any type in the high dose group of the 24-month study was 39 of 50 rats (78%) ( $p = 0.024$  for comparison with the control incidence of 22/54 rats [41%]; two-sided Fisher's exact test). Twenty-one of the 31 rats (68%) that died after the 12-month time-point (including the four that survived until the end of study) had prostatic inflammation, which was suppurative in nature in 19 rats and of the chronic type in two rats.

In the Levine et al. (1983) study, there was a shift from chronic inflammation to suppurative inflammation in the prostate with increasing RDX dose at 1.5 mg/kg-day and higher. This shift is statistically significant if tested using a Chi-squared test, with categories set for no lesions, chronic inflammation, and suppurative inflammation across all treatment groups ( $P < 0.0001$ ); prostatic inflammation was scored by Levine et al. as either chronic or suppurative. The shift was almost complete in the 31 rats that died in the 40 mg/kg-day group after 12 months on study, as only two animals had (minimal) chronic inflammation and 18 had suppurative inflammation. Although this analysis ideally should have taken into account mortality differences, the Levine study does not contain data that allows one to do this, as pointed out in the draft assessment.

- (c) The description of the methods used for histopathological evaluations lacked detail in Levine et al. (1983), which is an important issue given the large variation known for inflammation among the four prostate lobes, based on NTP data of aged F344 rats. The fact that the incidence of prostatic inflammation in the control group of the Levine *et al.* study was 40% lower than the range of inflammation incidences found in the dorsolateral prostate by Suwa et al. (2001) would suggest that some or many of the prostates examined by Levine et al. were ventral lobes, which have a low inflammation incidence (4-12%), according to Suwa et al. (2001). Suwa et al. indicated that there was considerable variation in which lobes were present and examined in the NTP studies they reviewed, suggesting that some of the study-to-study variation in the incidence of prostatic inflammation may be due to variations in which prostate lobes were examined.
- (d) There was no peer review or pathology working group review of the Levine et al. (1983) renal, bladder, and prostate pathology data, as was done for the liver lesions in female mice in Lish et al. (1984).
- (e) There may have been potential effects of the high prevalence of fighting among male rats in the highest dose (40 mg/kg-day) group and the resultant individual housing of these animals in the Levine et al. (1983) study. There is evidence in the literature that fighting may cause urogenital infections in male rats (Creasy et al. 2012). Because of this fighting, all males in the highest dose group were individually housed from 30-40 weeks into the study, which introduced another factor different from the other treatment groups that may have affected the animals in the 40 mg/kg-day group in uncontrolled ways.

In conclusion, the SAB found that the weight-of-evidence for identifying the prostate as a hazard of RDX exposure is sufficient because: (1) the Levine et al. (1983) study was considered sufficiently rigorous and appropriate for the time in which it was conducted and adequate to support the conclusion, even though the study has some deficiencies compared to current standards, and (2) the effects on the prostate were dose-related and statistically significant, albeit limited to the rat. The prostatic endpoint of all types of inflammation combined was not changed with increasing RDX dose, except at the highest dose (40 mg/kg-day), where its incidence was significantly increased. Only the incidence of suppurative inflammation and the shift from chronic to suppurative prostatitis were significantly increased at lower dose levels (1.5 mg/kg-day and higher). Both of the latter endpoints would be appropriate for analysis, but the SAB

agrees that the suppurative prostatitis incidence data is the most appropriate endpoint for quantitative risk assessment based on dose-response data.

### **Key Recommendations**

- Suppurative prostatitis should not be used as a surrogate marker of renal and overall urogenital effects, and instead, be considered a separate hazard of RDX exposure (see also Section 3.3.2.5.) and the appropriate prostatic endpoint for quantitative risk assessment.
- The description and analysis of prostatitis should be expanded to include discussion of both chronic and suppurative inflammation.
- The description of the various uncertainties regarding the Levine et al. (1983) rat study should be expanded to include commentary on lack of detail on methods used in histopathological evaluations, lack of peer review, and the impact of the high prevalence of fighting in highest dose rats.

### **3.3.2.2. Kidney and other urogenital system-specific toxicity values (Section 2.1.1).**

*Charge Question 3.b(ii). Is the selection of the Levine et al. (1983) study that describes kidney and other urogenital system effects scientifically supported and clearly described?*

The selection of the Levine et al. (1983) study that found kidney and other urogenital system effects is clearly described, but not entirely supported scientifically.

While the renal and bladder effects found in male rats in the high dose group of the study by Levine et al. (1983) were treatment-related and were the most likely cause of mortality in this group, the effects on the prostate were less straightforward [see also response to charge question 3b(v)].

One male in the lowest dose group (0.3 mg/kg-day) of 55 rats had renal papillary necrosis, but no other animals in the control or 1.5 and 8.0 mg/kg-day RDX dose groups had this lesion. By contrast, renal papillary necrosis was found in 33 of 50 male animals in the high dose group (40 mg/kg-day) (15 of 19 rats that died before the 12-month interim time point and 18 of 27 rats that died between 12 and 24 months – the lesion was not found in the four survivors at 24 months). Hemorrhagic/suppurative cystitis was found in 35 of 50 male rats of the high dose group, but in only one or two males per group in the lower dose groups and in none of the controls. These renal and bladder lesions tended to be more severe after 12 months of study than in rats examined at the six- and 12-month interim necropsies. Prostatic effects, namely a significant shift from chronic to suppurative inflammation, were seen at doses of 1.5 mg/kg-day and above and the overall incidence of prostatic inflammation was significantly increased in male rats in the high dose group (40 mg/kg-day).

The agency should improve the draft assessment's description and analysis of renal effects observed in studies other than those reported by Levine et al. The Levine et al. (1983) study was not the only animal study that found effects on the kidney. Renal medullary mineralization was reported by Martin and Hart (1974) in three of four males and three of four female Cynomolgus monkeys in the highest dose group tested (10 mg/kg-day), but not at lower RDX doses or in



controls. Cortical tubular nephrosis was found in four of ten males and one of ten female B6C3F1 mice at a very high RDX dose of 320 mg/kg-day, while this lesion was not present in control male or female mice (Cholakis et al. 1980). Both studies were of 90-day duration and the apparent renal effects were minimal to moderate in severity and not, or only marginally statistically significant. Cholakis et al. (1980) did not find any renal lesions in male F344 rats and only minimal microcalculi (mineralization) in one of ten female rats exposed to 40 mg/kg-day RDX via the diet for 90 days. In a two-generation study by Cholakis et al. (1980), renal cortical cysts, but no other renal lesions, were found in both control and treated CD (Sprague Dawley) rats.

Another 90-day study in F344 rats used lower doses by gavage and found no evidence of any treatment-related renal effects in males, while minimal-to-mild microconcretions (mineralization) were found in four of ten females that were administered RDX at a dose of 15 mg/kg-day and in seven of ten control females (Crouse et al. 2006). Levine et al. (1981a) found frequent microconcretions (mineralization) in female, but not in male, F344 rats, administered RDX via the diet for 90 days; control rats were without evidence of a treatment related effect. Levine et al. (1981a) also found nephropathy in both sexes, the incidence of which was reduced in the highest dose group (100 mg/kg-day); this reduction was significant in males but not in females. No renal toxicity was found in a 90-day dog study with dietary RDX doses up to 10 mg/kg-day (Hart et al. 1974). The only report of a lesion in the prostate came from the 90-day study by Crouse et al. (2006) in F344 rats administered RDX by gavage at a dose of 15 mg/kg-day; one of eight males had mild subacute inflammation in the prostate, while no prostate lesions were found in ten controls. There were no prostate lesions in any of the other 90-day studies mentioned above. In the 24-month study by Lish et al. (1984), a high frequency of cytoplasmic vacuoles in the renal tubular epithelium, with minimal-to-mild severity, was observed in male B6C3F1 mice at the six, 12, and 24-month time points; the male control group was an exception with only a 10% incidence of these cytoplasmic vacuoles at the six-month interim time point. Female mice had a low incidence of this renal change and this alteration was not reported in any of the other studies mentioned above.

In aggregate, mild toxic effects of RDX exposure on the kidney were found in some species, but not others, and in some studies in both sexes but in other studies only in male or female animals. Of note, some of these effects (mineralization) occurred in a small study with non-human primates, whereas some rodent studies did not find evidence of renal toxicity. Only in the chronic study of Levine et al. (1983) were severe toxic effects on the kidney found and they only occurred in males at the highest dose (40 mg/kg-day); bladder toxicity also occurred in this treatment group, whereas effects on the prostate occurred at doses of 1.5 mg/kg-day and above.

The marked sex difference in the renal toxicity due to RDX exposure found in rats by Levine et al. (1983) is not discussed in the draft assessment. However, there is precedent for a toxic chemical causing renal papillary necrosis selectively in male, but not female, F344 rats (Neal et al. 2003) and several drugs are well known for sex-specificity in their ability to cause renal papillary necrosis (Bach and Nguyen, 1998; Brix, 2002).

## **Key Recommendations**

- Improve the discussion and analysis of renal effects observed in studies other than those reported by Levine et al. (1983).
- Include a brief discussion of the marked sex difference in the renal toxicity in rats due to RDX exposure.

### **3.3.2.3.Points of Departure for Kidney and Other Urogenital System Endpoints (Section 2.1.2)**

*Charge Question 3b(iii). Is the calculation of a POD for this study scientifically supported and clearly described? Is the calculation of the HED for this study scientifically supported and clearly described?*

The SAB strongly recommends that suppurative prostatitis not be regarded as a surrogate marker for kidney and other urogenital system endpoints because of the various uncertainties that are associated with the hazard [see responses to charge questions 3b(i) and 3b(ii)]. If suppurative prostatitis is considered as a stand-alone endpoint (as recommended by the SAB), separate from kidney and other urogenital system endpoints, the calculation of both the POD and HED are scientifically supported and clearly described.

EPA's BMDS software was used to fit ten dose-response models to the data from Levine et al. (1983), and all models provided reasonable fits according to standard goodness-of-fit measures. Using a BMR of 10%, corresponding estimated BMDs for the models ranged from 1.67 to 10.8 mg/kg-day, with associated BMDLs ranging from 0.469 to 8.58 mg/kg-day. BMDLs from the ten models differ by more than threefold, so the lowest BMDL was selected, consistent with EPA guidance. The selected log-probit model has an estimated BMD of 1.67 mg/kg-day, which is within the range of study doses, thus obviating any issues of inappropriate extrapolation. The suppurative prostatitis POD for rats was determined to be 0.469 mg/kg-day.

Three methods were used to calculate the HED corresponding to the BMDL— one based on allometric scaling ( $BW^{3/4}$ ), another based on equivalent RDX serum AUCs in rats and humans at steady state, and a third based on equivalent RDX maximum serum concentrations in rats and humans after dosing. The methods for these calculations are clearly explained. The quality of data used for PBPK modeling is variable with respect to toxicity, but the resulting HED appears appropriate, with preference given to the AUC-based derivation. The SAB finds that the alternative approach of allometric scaling would introduce too many uncertainties.

## **Key Recommendation**

- The SAB strongly recommends that suppurative prostatitis should be used as a stand-alone endpoint, separate from kidney and other urogenital system endpoints for calculation of the POD and HED.

### **3.3.2.4.Uncertainty Factors for Kidney and Other Urogenital System Endpoints**

*Charge Question 3b(iv). Is the application of uncertainty factors to the POD scientifically supported and clearly described?*

The draft assessment used suppurative prostatitis in a two-year study in male rats (Levine et al. 1983) as a surrogate marker for the entirety of observed adverse effects of RDX exposure on the kidney and urogenital system. BMDS models were used to fit the data from Levine et al. (1983) using a 10% benchmark response rate (BMR). The human equivalent dose (HED) for the POD was calculated based on three methods. UFs were then applied to the BMDL10 HED to derive an RfD specifically for the kidney and urogenital system.

The SAB recommends that separate RfDs be derived for the kidney and urogenital system, based on findings of renal papillary necrosis and associated renal inflammation, and for suppurative prostatitis. This distinction designates the male accessory sex glands as a separate organ system, and challenges EPA's selection of suppurative prostatitis as a surrogate marker for the entirety of adverse effects on the kidney and urogenital system. This recommendation is in keeping with the fact that there is no known mechanistic link between suppurative prostatitis and renal papillary necrosis or adverse effects on renal function. Therefore, the charge question regarding the application of UFs can only be answered at this time for suppurative prostatitis, since an RfD has not been derived for renal papillary necrosis. Thus, the comments on the application of UFs are only relevant for the RfD derived based on suppurative prostatitis.

#### Intraspecies Uncertainty Factor ( $UF_H$ )

EPA applied an intraspecies uncertainty factor of 10 to account for toxicokinetic and toxicodynamic variability in the human population, which is standard default risk assessment practice. The SAB agrees with the use of a  $UF_H$  of 10. (See response to the  $UF_H$  response under Section 3.3.1.4 of this SAB report).

#### Interspecies Uncertainty Factor ( $UF_A$ )

An interspecies uncertainty factor,  $UF_A$ , of 3 ( $10^{1/2} = 3.16$ , rounded to 3) was applied to the point of departure to account for the remaining toxicodynamic and residual toxicokinetic uncertainty not accounted for in the toxicokinetic modeling. This is standard risk assessment practice where an adequate toxicokinetic model was applied to derive a human equivalent dose, and available data are not sufficient to define quantitative toxicodynamic differences between species. The SAB agrees with the application of a  $UF_A$  of 3.

#### Subchronic to Chronic Uncertainty Factor ( $UF_S$ )

The draft assessment used a  $UF_S$  of 1 to extrapolate from a subchronic experimental exposure duration to chronic exposure. The Levine et al. (1983) study was a chronic duration exposure study, and thus no extrapolation factor is needed. This SAB agrees that this is appropriate.

#### LOAEL to NOAEL Uncertainty Factor ( $UF_L$ )

The  $UF_L$  is meant to account for uncertainties in extrapolating from a LOAEL to a NOAEL when estimating an RfD. A  $UF_L$  of 1 was applied because the BMDL was used as a point of departure. Thus, there is no need to extrapolate from a LOAEL to estimate a NOAEL. This is standard risk assessment practice, and the SAB agrees that this is appropriate.

#### Database Uncertainty Factor ( $UF_D$ )

The assessment applied a  $UF_D$  of 3 in developing an RfD for suppurative prostatitis. The draft assessment notes that additional studies on neurotoxicity may provide a more sensitive endpoint

to use as the basis of an RfD. Thus, a UF<sub>D</sub> of 3 was applied across all points of departure, regardless of endpoint. In evaluating the RfD based on neurotoxicity, the SAB strongly recommends using a UF<sub>D</sub> of 10 rather than 3 due to database limitations. This UF<sub>D</sub> would be relevant to an RfD for suppurative prostatitis, if such an RfD were to be the basis of the overall RfD. However, if the RfD for suppurative prostatitis was only to be used specifically in a hazard index approach for this target, then an organ-specific UF<sub>D</sub> may be appropriate.

### **Key Recommendations**

- Develop or cite documentation for the use of organ-specific reference values for individual chemicals, including how these would be used in assessing the combined noncancer health impacts of multiple agents acting at a common site in cumulative risk assessments.
- A separate RfD should be derived for renal papillary necrosis and the associated renal inflammation of the kidney and urogenital system.
- The male accessory sex glands should be designated as a separate organ system with a separate RfD derived for suppurative prostatitis.

#### **3.3.2.5. Kidney and other urogenital system-specific reference dose (Section 2.1.4).**

*Charge Question 3.b.v. Is the organ/system-specific reference dose derived for kidney and other urogenital system effects scientifically supported and clearly characterized?*

The selection of suppurative inflammation of the prostate observed in the Levine *et al.* (1983) study in the draft assessment as a “surrogate marker” of the observed renal and urogenital system effects is not justified [see response to Charge Question 3b(i)] for derivation of a system-specific reference dose (RfD). Therefore, the organ/system-specific reference dose derived for kidney and other urogenital system effects is not sufficiently supported scientifically or clearly characterized.

### **Key Recommendations**

- A separate RfDs should be derived for renal papillary necrosis and the associated renal inflammation and for suppurative prostatitis.
- Available data are not consistent enough across species, doses, sex, or time points to recommend separate candidate RfDs for other, milder renal effects (tubular nephrosis, epithelial vacuolization, and mineralization) found in subchronic studies in mice, rats, and monkeys.

#### **3.3.3. Developmental and Reproductive System Effects**

##### **3.3.3.1. Developmental and Reproductive System Hazard**

*Charge Question 3c(i). The draft assessment concludes that there is suggestive evidence of male reproductive effects associated with RDX exposure, based on evidence of testicular degeneration in male mice. The draft assessment did not draw any conclusions as to whether developmental effects are a human hazard of RDX exposure. Please comment on whether the available human, animal, and mechanistic studies support these decisions. Are other hazards to human reproductive and developmental outcome adequately addressed?*

The SAB's response to Charge Question 3c(i) is subdivided into three components:

*No Evidence of Male Reproductive Effects*

The SAB disagrees with the conclusion in the draft assessment that there is suggestive evidence of male reproductive effects associated with RDX exposure. As outlined in detail in the SAB response to Question 3c(ii), the available animal evidence is too weak to support this statement. In addition, there are several animal studies that did not find effects on the male reproductive system. There is no human evidence indicating male reproductive toxicity; no human studies have focused on this question and there were no incidental reports of reproductive effects following RDX exposures. Furthermore, the mechanisms of action of RDX do not provide reasons to expect male reproductive toxicity.

*Conclusions as to whether Developmental Effects are a Human Hazard of RDX Exposure*

The SAB concludes that there are sufficient available data to draw the conclusion that RDX does not pose a risk of induction of structural malformations during human fetal development based on studies on rats and rabbits at doses that were high enough to occasionally produce maternal toxicity. The SAB encourages EPA to include this important piece of information for the overall risk assessment in the draft assessment. Additionally, the SAB agrees that conclusions cannot be drawn regarding other forms of developmental toxicity (decreased fetal weight, increased post-implantation loss), as these effects occurred only at maternally toxic dose levels.

The developmental toxicity observed was typical of findings associated with maternal toxicity and occurred at maternally toxic dose levels. It is generally understood that maternal toxicity, evidenced by body weight loss or reductions in body weight gain and/or decreases in food consumption, can contribute to developmental toxicity of the fetus in animal models. Developmental toxicity associated with maternal toxicity typically manifests as fetal weight reductions, increases in post-implantation loss (i.e., embryo/fetal death), and increases in the incidence of certain fetal skeletal variations. There is recognition within the scientific community of the possible effects on the fetus from maternal toxicity in common animal models [Carney and Kimmel, 2007; Rogers et al. 2005]. This concept was the primary topic discussed in an International Life Sciences Institute-Health and Environmental Sciences Institute (ILSI-HESI) sponsored working group, and the proceedings have been published [Beyer et al. 2011]. The findings in the RDX developmental toxicity studies of increased post-implantation loss, decreased fetal body weight and fetal skeletal variations, are those considered typically associated with maternal toxicity and occurred at maternal toxic dose levels.

In an embryo fetal developmental (EFD) toxicity study in F344 rats, maternal toxicity (mortality up to 31%) and developmental toxicity (reduced fetal body weights and increased resorptions) occurred at 20 mg/kg-day (Cholakakis et al. 1980). In Sprague Dawley rats administered 20 mg/kg-day RDX via gavage, maternal toxicity and increased resorptions were observed (Angerhofer et al. 1986). No structural malformations occurred at these doses or lower doses in either rat strain. Treatment in both of these studies starts on gestation day 6, while implantation is still in progress and ends on gestation day 15, prior to the closure of the hard palate. A longer dosage period as suggested for all current EPA (U.S. EPA, 1998a) and OECD (2015) guidelines, may have yielded more fetal toxicity, especially an effect on fetal weight.

The only two-generational study identified in the literature reported decreased offspring survival (including stillborn pups and postnatal deaths through the age of weaning) following a clearly maternally toxic dose of 50 mg/kg-day that was administered in the diet and adjusted approximately weekly (Cholakakis et al. 1980). Lower doses were not toxic to the dams or offspring.

Rabbits evaluated in an embryo-fetal developmental (EFD) toxicity study dosed on days 6 to 29 of gestation appear to be less sensitive than rats as exposures up to 20 mg/kg-day did not produce any maternal or embryo/fetal toxicity (Cholakakis et al. 1980). The rabbit embryo fetal development study at 0.2, 2 and 20 mg/kg-day showed fetal malformations with a low incidence at the 20 mg/kg-day dose and these changes were not present in control fetuses or seen at lower dose levels. The incidences ranged from 1 to 3 % and included a variety of malformations with no apparent biological relatedness, none of which were statistically significant. These data are difficult to put in context without a robust historical control database, systemic exposure levels and knowledge of the litter size. The report states that the findings were not statistically significant and thus not attributed to RDX exposure in rabbits.

#### Other Hazards to Human Reproductive and Developmental Outcome

Based on *in vitro* data, the SAB concludes that the mechanistic-based hazard demonstrating RDX inhibits GABAergic neurons presents a potential neurodevelopmental hazard that was not adequately addressed in the draft assessment. Several lines of evidence point to potential window(s) of susceptibility in the developing brain to chemicals interfering with GABAergic systems (see discussions in Sections 3.3.1.1, 3.3.1.2, and UFD discussion in Section 3.3.1.4).

A pilot developmental neurotoxicity study was conducted in rats that demonstrated significant accumulation of RDX in the immature brain of offspring and in the milk from dams treated with 6 mg/kg-day during gestation (Hess-Ruth, 2007). This dose level induced convulsions in adult animals. There were approximately equal concentrations ( $\mu\text{g/mL}$ ) in maternal blood and milk, and higher levels in younger postnatal day (PND) 1 pup brains compared to PND 10. A stated conclusion from this report was that further studies evaluating neurotoxicity and developmental effects of RDX should be conducted. It does not appear that a follow up study was conducted, thus no definitive assessment of potential developmental neurotoxicity in rats can be completed to inform risk for humans. Regardless, the SAB encourages the inclusion of a description of the potential mechanistic-based hazard in the draft assessment based on the reported mechanism to inhibit GABAergic neurons and the knowledge that RDX is present in the brain of developing rats during gestation and lactation.

#### **Key Recommendation**

- Due to significant weaknesses of the findings in the rat and mouse studies, RDX should not be considered a male reproductive toxicant.
- Animal studies do not suggest concerns for RDX causing fetal structural malformations to humans.

#### **Future Needs**

- Further studies should be performed to test the possibility that RDX has the potential to be a developmental neurotoxin (also noted in future needs in Section 3.3.1.4).

### 3.3.3.2. Reproductive System-Specific Toxicity Values

*Charge Question 3c(ii). Is the selection of the Lish et al. (1984) study that describes male reproductive system effects scientifically supported and clearly described?*

In consideration of all evidence, the SAB does not agree that the selection of Lish et al. (1984) for male reproductive effects is supported scientifically, and offers further suggestions on how to describe the data.

The SAB finds that the suggestive evidence of male reproductive effects provided by Lish et al. (1984), based on testicular degeneration in male mice exposed to RDX in their diet for 24 months, is weak, unsupported by other endpoints in that study showing no effect, complicated by the age of the mice and general toxicity of the RDX dose, and contradicted by most other studies.

In the study of Lish et al. (1984), the 10% and 11% incidence of testicular degeneration observed at doses of 35 and 108 mg/kg-day was not considered to be statistically significant. Also, no histological changes were observed at six or 12 months of study, durations that were much longer than the 1.4-month duration of spermatogenesis in mice. Furthermore, significant decreases in testis weight, which should have been observed if there were appreciable degeneration, were not observed.

The validity of 24-month chronic toxicity studies to evaluate male reproductive toxicity in rodents is questionable because of the loss of testicular function that occurs with aging in both rats and mice. In rats, the manifestations of aging in 2-year old animals include a high incidence of interstitial cell tumors (Cohen et al. 1978), declines in sperm production (Wang et al. 1993; Johnson & Neaves, 1983), reduced gonadotropin levels (Bruni et al. 1977), and reduced testosterone production due to aging of Leydig cells (Chen et al. 2002). In 2-year old mice, reductions in sperm counts and hormone levels were also observed (Bronson & Desjardins, 1977; Gosden et al. 1982), along with reductions in the numbers of stem spermatogonia, the loss of functional ability of these stem cells, and failure of the somatic environment to support spermatogonial differentiation (Suzuki & Withers, 1978; Zhang et al. 2006). Effects observed in rodents exposed to a potential reproductive toxicant in a 2-year chronic toxicity study might represent the combined effects of toxicant and aging, and not the result of prolonged treatment.

In addition, the indication of an effect of RDX on spermatogenesis suggested by Lish et al. (1984) is generally not supported by other studies (Table 3). In particular, Cholakis et al. (1980), using the same mouse strain, did not find any significant effects of RDX doses up to 320 mg/kg-day in a 3-month subchronic study. Although the RDX used by Cholakis et al. was of larger particle size than that used by Lish et al. which could reduce the uptake of RDX, mortality of the animals in the Cholakis et al. study administered 320 mg/kg-day was equivalent to that observed by Lish et al. at 175 mg/kg-day, indicating effective uptake of the RDX particles. Since 3-months allows for more than two complete rounds of spermatogenic cell differentiation, this should have been sufficient time to detect a toxic effect.

Furthermore, studies in rats indicate little male reproductive toxicity of RDX. In a 2-year chronic study, Hart et al. (1976) found no testicular degeneration or weight loss at doses up to 10

1 mg/kg-day. Similarly, Levine et al. (1983) found no effects of a dose of 8 mg/kg-day. However,  
2 at 40 mg/kg-day there was a significant decline in testis weight (14%) and a significant increase  
3 in the percentage of testes showing germ cell degeneration at 12 months of treatment. Although  
4 the effect was significant, the fact that there was 30% excess mortality by this time may indicate  
5 that the testicular damage was secondary to general toxicity. Data obtained at 24 months were  
6 not meaningful since all rats of this strain developed Leydig cell hyperplasia/neoplasms by 2  
7 years of age.

8  
9 Three 13-week subchronic studies in rats also failed to indicate significant testicular damage.  
10 Levine et al. (1981a, b; 1990) found no significant testicular effects of exposure at doses up to  
11 100 mg/kg-day. Also, Cholakis et al. (1980) found no changes in absolute testis weights or  
12 histopathological damage to testes at 28 or 40 mg/kg-day. Similarly, Crouse et al. (2006), in the  
13 only study using gavage, which had greater potency than dietary administration as indicated by  
14 20-30% mortality at doses of 10-15 mg/kg-day, reported no significant histological effects in  
15 testes or changes in absolute testis weights. The additional data of Cholakis et al. (1980) obtained  
16 as part of a 2-generational study, did indicate an 18% reduction in proportions of females  
17 impregnated by males exposed to RDX at 50 mg/kg-day. While this could reflect a testicular  
18 effect, it could also be a behavioral effect or a systemic effect, as suggested by the 14% excess  
19 mortality in this group.



**Table 3. Summary of Results of 7 Studies of Male Reproductive Toxicity of RDX**

Study	Species	Route	Significant Effect	Doses (mg/kg-day) Time (months)	Caveats	Negative Results (non-significant considered as negative)
Lish et al. (1984)	Mouse	Diet	None	35 & 108 24 mo.	Mortality <input type="checkbox"/> (>14%) Age-related effect	No histological change at 6 or 12 mo. The 10-11% incidence in testicular degeneration at 24 mo. was not significant. No decrease in testis weight
Cholakis et al. (1980)	Mouse	Diet	None	40, 80, 160, 320 3 mo.		No histological changes No decrease in testis weight
Levine et al. (1983)	Rat	Diet	40% increase in incidence of germ cell degeneration 14% decrease in testis weight	40 mg/kg- day 12 mo.	Mortality <input type="checkbox"/> 31% at 12 mo.	No effects at 8 mg/kg-day No effects at 6 months with 40 mg/kg-day No germ cell degeneration at 40 mg/kg-day at 24 mo.
Hart (1976)	Rat	Diet	None	10 mg/kg- day 12 & 24 mo.		No histological changes (24 months) No decrease in testis weight
Cholakis et al. (1980)	Rat	Diet	18% reduction in proportion of females impregnated †	50 mg/kg- day 3 mo.	Mortality <input type="checkbox"/> 14% Possible behavioral effect	Reduction in impregnation not observed at 16 mg/kg- day No histological changes at 40 mg/kg-day No decreases in testis weight at 28 or 40 mg/kg-day
Levine (1981a,b; 1990)	Rat	Diet	None	10, 30, 100 3 mo.		No histological changes No decreases in testis weight
Crouse et al. (2006)	Rat	Gavage	None	15 mg/kg- day 3 mo.		No histological changes No decreases in testis weight

☐ Excess mortality compared to observed in controls

† Calculated as significant by reviewer using Chi-square test at P=0.004

Finally, the SAB did not find the selection of Lish et al. (1984) to be clearly described, and provided specific comments in Appendix B on the text, tables and figures to improve presentation of data on reproductive and developmental toxicity.

**Key Recommendation**

- SAB finds that male reproductive toxicity should not be used for derivation of a reproductive-system specific toxicity value, as there have been no convincing studies showing significant male reproductive toxicity.

**3.3.3.3.Points of Departure for Reproductive System Endpoints**

*Charge Question 3c(iii). Is the calculation of a POD for this study scientifically supported and clearly described? Is the calculation of the HED for this study scientifically supported and clearly described?*

As discussed elsewhere [e.g., in response to questions 3c(i) and 3c(ii)], the SAB does not support use of the Lish et al. (1984) study for describing male reproductive system effects, because the suggestive evidence of male reproductive effects provided by Lish et al. (1984) is weak, unsupported by other endpoints in that study showing no effect, complicated by the age of the mice and general toxicity of the RDX dose, and contradicted by most other studies. Given that Lish et al. (1984) was the data source for dose-response modeling and subsequent derivations of the POD and HED, these derived POD and HED are not scientifically supportable.

**Key Recommendation**

No POD for reproductive endpoints should be calculated from the existing data and therefore there is no need to calculate the HED.

**3.3.3.4.Uncertainty Factors for Reproductive System Endpoints.**

*Charge Question 3c(iv). Is the application of uncertainty factors to the POD scientifically supported and clearly described?*

The draft assessment used the data on testicular degeneration in mice from a 2-year dietary study (Lish et al. 1984) as the basis for derivation of the POD. BMDS models were used to fit the incidence data to derive a BMDL for a 10% BMR. Three methods were used to derive an HED from the mouse POD. The draft assessment notes that the toxicokinetic data available for the mouse are not as robust as for the rat, and thus confidence in the use of PBPK modeling to account for interspecies toxicokinetics is low. Rather, the default allometric scaling approach was used to derive the HED by scaling dose by  $\frac{3}{4}$  power of body weight. After adjusting the mouse POD to an HED with this scaling, UFs were applied to derive an RfD for male reproductive toxicity.

Since SAB finds that the suggestive evidence of male reproductive effects cited by EPA in Lish et al. (1984) is not scientifically defensible, the SAB does not support derivation of an RfD based on male reproductive system effect [see response to Charge Question 3c(iii)], and concludes that an RfD based on testicular degeneration is not supported scientifically. The question of UFs as applied to the POD is therefore extraneous.

**Key Recommendation**

- Since no valid POD should be calculated for reproductive endpoints, there is no need to discuss UFs for reproductive endpoints.

**3.3.3.5.Reproductive System-specific Reference Dose.**

*Charge Question 3c(v). Is the organ/system-specific reference dose derived for reproductive system effects scientifically supported and clearly characterized?*

The RfD for reproductive effects based on testicular degeneration is clearly described but not scientifically supported. Reasons for this conclusion are provided above in response to question 3c(ii), and briefly summarized below.

Testicular degeneration was reported at terminal sacrifice (24 months) in one 2-year dietary study in mice (Lish et al. 1984). Germ cell degeneration was also observed in a 2-year dietary study in rats (Levine et al. 1983) but only at the 12-month interim sacrifice and not at the 6-month interim or 24-month terminal sacrifice. The SAB notes that testicular histopathology should have been seen at earlier time points (e.g., the 6-month and 12-month interim sacrifices) in Lish et al. (1984), as these exposure durations were several times longer than the 1.4-month duration of spermatogenesis in mice. Further testicular degeneration was not observed in the majority of the dietary and gavage studies in rodents (5 of 7 showed no effect).

Other reproductive effects observed included changes in testicular absolute and relative weight, but these findings were inconsistent across studies. Effects on fertility were noted in a 2-generation reproductive study in CD rats at the high dose (50 mg/kg-day) (Cholakakis et al, 1980), but both the male and female rats had decreased weight gain and increased mortality. Thus, it was difficult to attribute the reduction in fertility to a specific reproductive toxicity effect of RDX. In a dominant lethal assay (Chokakis et al. 1980), the decreased rates of pregnancy of untreated females that were mated with F0 males treated with RDX at 50 mg/kg-day may have been associated with generalized toxicity in the treated males rather than a specific effect of RDX. There were no observations of histological changes in the testis or decreased testicular weight in any of the treated animals in Cholakakis et al. (1980).

The Agency provided the BMDS analysis in Appendix D of the Supplemental Document, and clearly described the rationale for deriving the HED and applying the UFs. However, since the toxicological effect used as the basis of the RfD was testicular degeneration, and this is not supported scientifically, then the RfD is not supported scientifically.

**Key Recommendation**

- No RfD based on male reproductive toxicity should be calculated since no valid POD can be estimated.

**3.3.4.Other Noncancer Hazards**

*Charge Question 3d. The draft assessment did not draw any conclusions as to whether liver, ocular, musculoskeletal cardiovascular, immune, or gastrointestinal effects are human hazards*

1 *of RDX exposure. Please comment on whether the available human, animal and mechanistic*  
2 *studies support this decision. Are other non-cancer hazards adequately described?*  
3

#### 4 Liver, Ocular, Musculoskeletal, Cardiovascular, Immune, and Gastrointestinal Effects

5 In the process of identifying the health hazards of RDX, a conclusion should be made for each  
6 hazard endpoint discussed based on the available evidence streams (human, animal, and  
7 mechanistic), and a critical evaluation of the quality and relevance of the data reviewed. In this  
8 regard, the SAB recommends that the draft assessment be explicit as to whether or not the  
9 available evidence supports each of the discussed systemic effects as a potential human hazard,  
10 and the rationale for reaching that conclusion. Furthermore, the meaning of the statement (line  
11 11, p. 1-60 and lines 13 & 14, p. 1-69) “at this time no conclusions are drawn regarding [viz.  
12 liver effects or other non cancer effects] as human hazards of RDX exposure”, is not clear. Does  
13 the draft assessment mean to state that existing data are inadequate to establish that RDX can  
14 cause a particular adverse effect in humans, or that existing data are inadequate quantitatively to  
15 serve as a point of departure for a risk assessment? Clearly, very high unspecified doses of RDX  
16 cause modest, reversible increases in liver-specific serum enzyme activities in humans. High  
17 doses of RDX cause modest increases in serum enzyme activities and hepatomegaly in dogs, and  
18 while RDX does not appear to enhance serum enzymes in rats, it produces increased liver  
19 weight. Still, increases in relative liver weights are not observed consistently from one study to  
20 another. In light of this example, high doses of RDX clearly cause modest hepatic effects in  
21 humans and animals, though there are no dose-response data available to calculate an RfD.  
22 Simply stating that “no conclusions are drawn regarding liver effects as a human hazard of RDX  
23 exposure” leaves the reader uncertain as to what decision EPA has made and why.  
24

25 The description of Liver Effects in Section 1.2.4 is well written and comprehensive. The authors  
26 have done an excellent job grouping studies and providing detailed accounts. Conclusions about  
27 consistency of inter- and intra-species findings of different durations are scientifically  
28 appropriate. The integration of the liver effects on the top of the page 1-61 should lead to a more  
29 specific/definitive conclusion, as addressed above, rather than the conclusion stated in lines 10  
30 and 11.  
31

32 It is recommended that the overviews of ocular, cardiovascular (CV), musculoskeletal (MS),  
33 immune, gastrointestinal (GI) and hematological effects be moved from Section C.3.2 of  
34 Appendix C to a new subsection 1.2.5 (Other Noncancer Effects), rather than be included in an  
35 Appendix that readers may not be able to readily access. As such, section 1.2.5 (Carcinogenicity)  
36 would become section 1.2.6.  
37

38 The accounts of ocular, cardiovascular system, musculoskeletal system, immune system,  
39 gastrointestinal system, and hematological system effects of RDX, like those for the liver, are  
40 generally detailed, accurate and comprehensive in their coverage of each organ system. It is  
41 laudable that each account, with the exception of the musculoskeletal system, is concluded by a  
42 definitive, well-supported summary statement of the available evidence streams. However, these  
43 summaries lack a conclusion and rationale for whether the evidence supports potential human  
44 hazards from RDX exposure. The following additional information may be helpful in developing  
45 conclusions and related rationale.  
46

It is stated in lines 24-26 of page C-44 that muscle injury was indicated by elevated levels of AST or myoglobinuria. However, some other enzymes measured in serum are more specific for muscle damage. Kucukardali et al. (2003), for example, reported transient increases in several serum enzymes in four of five patients experiencing RDX-induced seizures. One of the enzymes, creatine phosphokinase (CPK), is primarily indicative of muscle damage. Testud et al. (2006) measured elevated CPK and myoglobin levels in an Octogen-poisoned patient. The clinicians attributed these findings to muscle damage secondary to seizures.

The EPA concludes in lines 5 and 6 of page C-46 that histopathological changes have generally not been reported in RDX dietary studies. Kucukardali et al. (2003) did observe gastroduodenitis by endoscopy in three of five human poisoning victims who ingested enough RDX to cause protracted seizures. Severe irritation of the gastrointestinal mucosa by direct contact with RDX would account for the nausea and vomiting commonly experienced by humans who ingest high doses of the chemical.

With respect to effects of RDX on the immune system, the empirical data have been summarized adequately in Appendix C.3.2. Based on the available animal studies, consistent dose-related immune system effects from oral exposure to RDX were not observed. However, it should be noted that none of these studies, including that of Crouse et al. (2006), completed sensitive immune function evaluations. The Crouse study was specifically designed to evaluate immunotoxicity in rats, but included only less sensitive structural evaluations of the immune system, such as populations of red and white blood cells, proportion of cell surface markers, cellularity in proportion to organ weight, B and T cells in the spleen, and CD4/CD8 antigens of maturing lymphocytes in the thymus). As observed by USEPA (1998), WHO (2012), and others, evaluation of such structural parameters in the absence of more sensitive functional testing is unlikely to detect immunosuppression, unintended immune stimulation, autoimmunity, or dysregulated inflammation.

#### Other Non-Cancer Hazards:

The potential “other non-cancer hazards” from RDX exposure are identified and discussed in Section 1.2.4 and 1.3.1 (liver), and Section 1.2.6 and Appendix C.3.2 (ocular, musculoskeletal, cardiovascular, immune system, gastrointestinal, and hematological) of the draft assessment. In Appendix C.3.2, lines 5-6 it states “Overall, at the present time, the evidence does not support identifying these other systemic effects as human hazards of RDX exposure.” In the subsequent paragraphs summarizing the evidence for the other systemic effects, the text does not provide a clear rationale for why the evidence does not support the listed effects as potential human hazards. For example, is it due to insufficient data, inconsistent data, or sufficient data to conclude that these health endpoints are not sensitive endpoints?

Importantly, neuroinflammation has emerged as a key characteristic of most neurological conditions, including seizure and epilepsy, as recently reviewed by Dey et al. (2016) and Eyo et al. (2017). In particular, RDX-induced seizures may trigger acute immune and inflammatory responses within the brain, while chronic neuroinflammation may result from recurrent seizures. Neuroinflammation, in turn, has a proconvulsant effect by lowering the seizure threshold, influencing seizure severity and recurrence. This context is relevant to the interpretation of studies in which RDX exposures provoked convulsions. It is less clear what relationship, if any,

there may be between less severe manifestations of RDX neurotoxicity and neuroinflammatory or other chronic immune system responses.

Not addressed in the draft assessment were the dose-related effects on body weights and/or body weight gains, although this was identified as a potential adverse effect of RDX elsewhere (e.g., Sweeney et al. 2012a, b; EPA, 2012b). Dose-related decreases in body weight gain were frequently observed in repeated dose studies with RDX, and should also be considered and discussed as a potential noncancer effect. Reduction in body weight is a common manifestation of adverse effects of chemicals, reflecting generalized systemic toxicity. This parameter has been utilized in numerous IRIS assessments for the derivation of reference values. The RDX literature should be screened to identify subchronic or chronic animal studies in which dose-dependent decreases in body weight/body weight gain are reported. Dose-related body weight effects should be discussed in the draft assessment, including their suitability to carry forward from hazard identification to the dose-response analysis.

### **Key Recommendations**

- For each of the other noncancer hazards discussed in the draft assessment, add a summary statement regarding whether the available studies do or do not support a conclusion that the identified toxicity is a potential human hazard. Include an explanation of the rationale for reaching the conclusion, taking into consideration the additional information pertaining to liver effects, the muscle injury, immune system, neuroinflammation and gastrointestinal effects, as detailed above by the SAB.
- Include as a potential noncancer hazard the available subchronic and chronic data on body weight/body weight gain, and whether the studies do or do not support a conclusion that body weight effects represent a potential systemic human hazard. Discuss the rationale for the conclusion and explain why body weight effects are or are not carried forward to the dose-response analysis.

### **Suggested Recommendation**

- Move the overviews of ocular, cardiovascular, musculoskeletal, immune, gastrointestinal and hematological effects from Section C.3.2 of the Appendix C to Section 1.2 of the main body of the draft assessment. These overviews should be placed in subsection 1.2.5 (Other Noncancer Effects) rather than be part of an Appendix. Section 1.2.5 (Carcinogenicity) would become section 1.2.6.

## **Cancer**

### **3.3.5.1. Cancer Hazard**

*Charge Question 3e(i). There are plausible scientific arguments for more than one hazard descriptor as discussed in Section 1.3.2. The draft assessment concludes that there is suggestive evidence of carcinogenic potential for RDX, and that this descriptor applies to all routes of human exposure. Please comment on whether the available human, animal, and mechanistic studies support these conclusions.*

1 The SAB concurs with the EPA that “*suggestive evidence of carcinogenic potential*” is the most  
2 appropriate cancer hazard descriptor for RDX and that this descriptor applies to all routes of  
3 human exposure

4  
5 In the draft assessment, the EPA considered two potential hazard descriptors under the EPA’s  
6 *Guidelines for Carcinogenic Risk Assessment* (USEPA, 2005): “*likely to be carcinogenic to*  
7 *humans*” and “*suggestive evidence of carcinogenic potential*,” with the latter indicative of a  
8 lesser weight of evidence. Per the guidelines, the *suggestive evidence of carcinogenic potential*  
9 descriptor is “appropriate when the weight of evidence is suggestive of carcinogenicity, a  
10 concern for potential carcinogenic effects in humans is raised, but the data are judged not  
11 sufficient for a stronger conclusion.” A *likely to be carcinogenic in humans* descriptor is  
12 appropriate when “the weight of evidence is adequate to demonstrate carcinogenic potential to  
13 humans” but is not strong enough to justify the highest weight of evidence descriptor  
14 *carcinogenic in humans*.

15  
16 The draft assessment noted that RDX tested positive in more than one species, sex, and strain in  
17 animal studies and that this evidence was consistent with a “*likely to be carcinogenic in humans*”  
18 descriptor, as provided in the EPA guidelines (USEPA, 2005), and suggested that more than one  
19 descriptor might apply to RDX. However, the draft assessment also noted that the evidence of  
20 carcinogenicity outside the B6C3F1 mouse was not robust, and this factor was decisive in  
21 choosing a hazard descriptor, which was “*suggestive evidence of carcinogenic potential*”.

22  
23 In considering the most appropriate cancer hazard descriptor for RDX, the SAB evaluated the  
24 strength of evidence for positive cancer findings. The SAB agrees with the EPA that the relevant  
25 observations are the liver tumors that were observed in female B6C3F1 mice and male F344 rats  
26 and lung tumors that were observed in female B6C3F1 mice in two-year dietary bioassays (Lish  
27 et al. 1984; Levine et al. 1983) and identified other limitations that raised concerns.

28  
29 The findings of the SAB are as follows:

- 30  
31 1) *Mortality in the high dose groups*. The high dose in the Lish et al. dietary study in mice was  
32 initially 175 mg/kg; however, the dose of RDX was reduced at week 11 of the study to 100  
33 mg/kg diet, due to acute toxicity and high early mortality (30 of 65 males and 36 of 65  
34 females). The acute toxicity and early mortality reduced the effective number of animals after  
35 11 weeks on the study to 35 male mice and 29 female mice. Twenty-two male and 25 female  
36 mice in the high dose group survived to the scheduled sacrifice at 24 months. Similarly, the  
37 high dose in the Levine et al. dietary study in rats was 40 mg/kg-day, and mortality was high  
38 throughout the study period of 24 months. Unlike the mouse study, mortality occurred  
39 gradually over the entire period of the rat study, with mortality in most rats occurring after 6  
40 months. Male rats were particularly affected by RDX toxicity, and histopathological  
41 evaluations indicated that the high mortality was likely due to renal disease. Four of 55 males  
42 and 28 of 55 females in the 40 mg/kg dietary exposure survived to scheduled sacrifice. The  
43 *Guidelines for Carcinogen Risk Assessment* states that “If toxicity or mortality is excessive at  
44 the high dose, interpretation [of cancer] depends on whether or not tumors are found. ...  
45 Studies that show tumors at lower doses, even though the high dose is excessive and may be  
46 discounted, should be evaluated on their own merits.”

- 1  
2 2) Liver tumors in rats. A positive finding of cancer hazard in two species is based on the liver  
3 tumor response in male F344 rats to RDX, in addition to the liver and lung tumors observed in  
4 female RDX-exposed mice. The liver tumor response of males to RDX in the Levine et al.  
5 (1983) rat study was only significant in a trend test, if the incidence of hepatocellular  
6 carcinomas in males of the high dose group (40 mg/kg-day) was included. When this group  
7 was omitted from the analysis, because of its high mortality, the trend was not significant nor  
8 was a pair-wise comparison of the high dose group incidence to that of control males. There  
9 was no dose-related trend for the incidence of adenomas or the combination of adenomas and  
10 carcinomas. Although the incidence of benign liver tumors in control males was on the high  
11 end of the range for historical controls (Haseman et al. 1985), the evidence for an association  
12 of RDX exposure with increased liver tumors in this rat study is weak.  
13
- 14 3) Lung tumors in mice. The increased incidences of liver and lung tumors observed in female  
15 B6C3F<sub>1</sub> mice would support a positive finding of cancer hazard at two sites. However, the  
16 lung tumor response to RDX in the mouse study of Lish et al. (1984) showed a significant  
17 trend for an increase only when the incidence of lung tumors (adenomas and carcinomas  
18 combined) in females of the high dose group (175/100 mg/kg-day) was included in the trend  
19 test. The trend and pairwise comparison tests were not significant if the high dose group was  
20 excluded from the analysis. A positive trend for the incidence of pulmonary carcinomas (not  
21 adenomas) was observed in both sexes of mice, but only when the high dose groups were  
22 included. The incidence of these tumors was quite close to that observed in historical controls  
23 (Haseman et al. 1985). Thus, the evidence for an association of RDX exposure with increased  
24 lung tumors in this mouse study is weak and solely driven by the findings in the high dose  
25 group that suffered from high early mortality.  
26
- 27 4) Liver tumors in mice. A positive finding of cancer hazard in both sexes of one species is  
28 based on the liver tumor response in male and female B6C3F<sub>1</sub> mice to RDX, but the SAB  
29 identified several concerns regarding the liver tumors in mice.
- 30 • Although there were suggestive increases in liver tumors in male mice, none of the  
31 increases appear to be statistically significant, using either a trend test or pairwise  
32 comparison tests. Of note, the incidences of these tumor types are quite variable in mice  
33 and the observed increases are within the range of incidences observed in historical  
34 controls (Haseman et al. 1985). Of further note is that the incidence of combined adenoma  
35 and carcinoma liver tumors observed in the high dose group is near the high end of the  
36 historical control range and that increases in tumors at other sites were not observed in  
37 male mice.
  - 38 • The liver tumor findings were more robust in female mice, but there were also concerns  
39 with these observations, due to the unusually low incidence of hepatocellular tumors in  
40 female control mice. None of the concurrent female mice controls had hepatocellular  
41 carcinomas (0.0%) and one of 65 had a liver adenoma (1.5%), while historical incidence  
42 control data published by the NTP were 8.0% (range 0-20%) for the combined  
43 hepatocellular carcinoma and/or adenoma, indicating that the observed 1.5% incidence was  
44 notably at the low end of the range of incidences found in historical controls.
  - 45 • In a reevaluation of hepatocellular neoplasms in female mice by a Pathology Working  
46 Group (PWG), the original histological sections from female mice were retrieved and a



second examination was performed by pathologists (Parker et al. 2006). It was noted that a reevaluation of neoplasm sections from just one sex is unusual; sections from both male and female animals would be reevaluated typically to ensure that findings in both sexes were reliable. Members of the PWG then reexamined all hepatocellular neoplasm sections from female mice and cited factors that reduced their confidence in a positive interpretation of the study. The cited factors included variations in the number of liver sections per mouse, the absence of precursor lesions, such as foci of cellular alteration, and, most importantly, the low incidence of hepatocellular neoplasms (1.5%) in the control females. A discrepancy in the number of mice examined by Lish et al. (1984) and by Parker et al. (2006), while not major, further undermined the confidence in the quality of the data.

- 5) Non-neoplastic histopathological liver changes. Non-neoplastic histopathological changes in the liver were absent in the majority of subchronic studies available in the literature, and pre-neoplastic lesions were absent in the livers of mice and rats at interim sacrifices conducted at 6 and 12 months in the two-year bioassays by Lish et al. (1984) and Levine et al. (1983). The finding that non-neoplastic changes in livers were not associated with RDX exposure in the majority of animal studies suggested that intrinsic factors may be involved in the observed tumor findings, especially in light of the fact that the mode of action of RDX carcinogenicity cannot be determined based on the current understanding of RDX metabolism (see below). It is acknowledged that the absence of hepatic precursor lesions does not, in itself, negate the possibility that RDX could have caused the increases in liver neoplasms, but nevertheless is a weight of evidence factor to consider.
- 6) The lack of pathology peer-review and available data to support mortality-based statistics for neoplasms in the two-year bioassays by Lish et al. and Levine et al. Carcinogenicity findings in well-conducted experimental animal studies are regarded as evidence of potential cancer risk to humans by national and international health agencies. In order for experimental animal studies to serve as reliable sources of data for the evaluation of the carcinogenic potential of environmental agents, certain criteria should be met (Melnick et al. 2008). These include: a) animal models that are sensitive to the end points under investigation; b) detailed characterization of the agent and the administered doses; c) challenging doses and durations of exposure (approximately 2 years for rats and mice); d) sufficient numbers of animals per dose group to be capable of detecting a true effect; e) multiple dose groups to allow characterization of the dose-response relationships; f) complete and peer-reviewed histopathologic evaluations; and g) pairwise comparisons and analyses of trends based on survival-adjusted tumor incidence (Melnick et al. 2008). The Lish et al. and Levine et al. studies met criteria a – e; however, complete and peer-reviewed histopathologic evaluations and pairwise comparisons and analyses of trends based on survival-adjusted tumor incidence were not conducted and the available data did not allow EPA to perform these analyses. Additionally, necropsy and histological processing records were not available to link gross lesions observed at necropsy or the number of gross lesions with histological sections that were evaluated for each animal.
- 7) Limited evidence to support a mode of action for RDX carcinogenicity. Data are not available in the literature to support the metabolism of RDX by human liver or lung enzymes or by human microflora to form genotoxic agents. One rodent study demonstrated the reductive

transformation of RDX to N-nitroso compounds (Pan et al. 2007b). It is unclear if this transformation occurred via microflora, non-enzymatic processes, or by rodent metabolic enzymes. Bhushan et al. (2003) reported that rabbit cytochrome P450 2B4 converts RDX to 4-nitro-2,4-diazabutanal *in vitro*. This compound and 4-nitro-2,4-diaza-butanamide were identified as minor end product metabolites in the urine of Yucatan miniature pigs (Major et al. 2003). However, the genotoxic potential of these compounds has not been determined in mutagenesis assays. Numerous studies have shown that RDX yields negative test results with the Ames *Salmonella typhimurium* assay in a variety of bacterial strains (Cholakakis et al. 1980; George et al. 2001; Tan et al. 1992) and is not cytotoxic or mutagenic in the *in vitro* mouse lymphoma test or *in vivo* by the mouse bone marrow micronucleus test (Reddy et al. 2005). RDX was reported by one group to be weakly mutagenic in one strain of *Salmonella typhimurium* using S9 activation (Pan et al. 2007a) and showed some evidence of mutagenic activity in *Vibrio fischeri* using the Mutatox assay (Arfsten et al. 1994). *In vitro* biotransformation studies on RDX suggest that RDX can be metabolized by anaerobic bacteria in soils to form N-nitroso derivatives. Such biotransformation has also been demonstrated in the mammalian gastrointestinal tract (Major et al. 2007; Musick et al. 2010; Pan et al. 2007b). These minor N-nitroso metabolites, MNX and TNX, were reported to be positive in *in vitro* genotoxicity studies using several strains of *Salmonella typhimurium* (Pan et al. 2007a; George et al. 2001). Moreover, MNX was reported to be positive in genotoxicity studies in mammalian cells *in vitro* with metabolic activation with S9 (Snodgrass, 1984). Other modes of action of RDX, such as oxidative stress mechanisms in carcinogenesis, have not been investigated. However, it should be noted that, while understanding the mode of action can sometimes support a concern for carcinogenicity of a chemical, it is not requisite to the determination of its cancer hazard.

Based on the guidance provided in the EPA's *Guidelines for Carcinogenic Risk Assessment* (USEPA, 2005) and points 2-4 above, the SAB considers that the evidence for a positive tumor response to RDX in two species, two sexes, or two sites required by EPA for a "*likely to be carcinogenic in humans*" descriptor is weak or absent. Given the limitations and nature of the carcinogenicity data available, the SAB concludes that the descriptor, "*suggestive evidence of carcinogenic potential*", is appropriate. As noted in the draft assessment and in the discussion above, oral exposure to RDX has been observed to result in tumors in liver, which is beyond the point of initial contact. This is indicative of carcinogenic effects that are systemic rather than confined to the portal of entry to the body, and thus carcinogenic potential is independent of the route of exposure. Therefore, the SAB agrees with EPA that this descriptor applies to all routes of exposure.

#### **Key Recommendation**

- Strengthen and make more specific the justification for selecting the "*suggestive evidence of carcinogenic potential*" descriptor rather than the "*likely to be carcinogenic to humans*" descriptor.

#### **Suggested Recommendations**

- Expand the discussion to include more on the limitations of the Lish et al. (1984) and Levine et al. (1983) animal studies.
  - Clarify that the absence of hepatic precursor lesions in the female mice of the Lish et al.

(1984) study does not, by itself, negate the possibility that RDX could have caused the increases in liver neoplasms.

- o Include a more complete description of the differences in mortality time course between mice in the Lish et al. (1984) study and rats in the Levine et al. (1983) study administered the high dose level of RDX in the diet and the potential impact of these differences on the interpretation of the hepatic and pulmonary neoplasms in female mice.

#### **3.3.5.2. Cancer-specific Toxicity Values.**

*Charge Question 3e(ii). As noted in EPA's 2005 Guidelines for Carcinogen Risk Assessment, "When there is suggestive evidence, the Agency generally would not attempt a dose-response assessment, as the nature of the data generally would not support one; however, when the evidence includes a well-conducted study, quantitative analyses may be useful for some purposes, for example, providing a sense of the magnitude and uncertainty of potential risks, ranking potential hazards, or setting research priorities." Does the draft assessment adequately explain the rationale for quantitative analysis, considering the uncertainty in the data and the suggestive nature of the weight of evidence, and is the selection of the Lish et al. (1984) study for this purpose scientifically supported and clearly described?*

The SAB finds that the draft assessment adequately explains the rationale for a quantitative analysis of RDX cancer assessment and that the selection of the Lish et al. (1984) study for this purpose is supported scientifically and clearly described.

Despite concerns associated with interpretation of the data as discussed in response to Charge Question 3e(i) (Section 3.3.5.1), the Lish et al. (1984) study was a well-conducted two-year bioassay that included a large number of animals tested at multiple dose levels, and increased incidences of neoplasms occurred in exposed female mice.

#### **Key Recommendation**

- The Lish et al. (1984) study is suitable and appropriate for dose-response assessment, consistent with the EPA's 2005 *Guidelines for Carcinogen Risk Assessment* (USEPA, 2005).

#### **3.3.5.3. Point of Departure (POD) for Cancer Endpoints.**

*Charge Question 3e(iii). Are the calculations of PODs and oral slope factors scientifically supported and clearly described?*

The SAB finds that the calculations in the draft assessment of the PODs and oral slope factors for cancer endpoints are not clearly described, and the SAB expresses the following concerns about whether these are scientifically supported. As described further below, the SAB expresses a number of concerns with the data used to derive the cancer POD, with the rationale for restricting modeling to the multistage model, and with the conditions under which the agency's MS-COMBO modeling methodology provides a valid POD and cancer slope factor estimate.

The draft assessment discusses two modes of action for cancer, genotoxicity and cell proliferation, but the available data are too limited to support use of a mechanism-based model.

Without a clear mode of action, the linear low-dose extrapolation method recommended in the EPA 2005 Cancer Guidelines is used in the draft assessment. The SAB agrees with this choice.

The SAB has concerns with the low incidence of liver tumors (hepatocellular adenomas and carcinoma) in female mice and its impact on dose-response modeling. As indicated in Section 1.2.5 of the draft assessment, the 1.5% incidence of liver tumors in the control B6C3F1 mice is unusually low. This was reported by the study authors as significantly lower than reported for historical controls, and at the low end of the range of the incidences in control females reported for this mouse strain by the National Toxicology Program (NTP) (mean 8%, range 0-20%). This low control incidence could significantly influence the estimate of the POD. The SAB recommends that for liver cancer, additional BMD modeling (i.e., a sensitivity analysis) be performed to examine and illustrate the impact of low concurrent controls on model choice and POD estimate.

The draft assessment discusses concerns that inclusion of the highest dose in the dose-response model for liver tumors for the B6C3F1 mouse study may impact the POD estimate and estimated cancer slope factor. The previous RDX risk assessment excluded the high dose used in this study in deriving the POD and cancer slope factor. A change in the highest dose at week 11 due to high mortality was reported, and mice that died prior to week 11 were excluded from the analysis. This results in a reduced sample size for the highest dose group from 65 to 31 animals and subsequent increased uncertainty in the response to this dose. While survival times of mice in the highest dose group were not significantly different from other dose groups, high mortality in the early weeks may mean that remaining survivors had other differences that potentially resulted in higher resistance to cancer. Excluding the highest dose group results in a fitted multistage model form that is almost linear. Using this fitted model produces an estimated POD that is much less than that estimated with the highest dose group included. This alternate POD in turn produces an unrealistically high cancer slope estimate (see Figure D-15). The SAB has no specific recommendation on how EPA should address this issue other than to include/exclude the highest dose in the sensitivity analysis for examining concurrent controls.

The draft assessment relies on the multistage model to describe dose-response relationships and subsequently to estimate the POD and cancer slope factor. As discussed in the BMD guidance document (USEPA, 2012a), the IRIS program prefers to use a multistage model for cancer dose-response modeling of cancer bioassay data, when no biological basis for model form is available because it considers this class of models as sufficiently flexible to address the typical dose-response patterns of cancer bioassay data, and its use encourages comparability across IRIS assessments. The draft assessment does not include a rationale for using the multistage model in the draft assessment, and this omission leads the SAB to question the validity and scientific adequacy of other aspects of the dose-response modeling, as well as the use of the MS-COMBO package. Furthermore, the SAB concludes that more discussion on the rationale for using the multistage model and how the EPA typically assesses the multistage model fit would greatly improve an understanding and reduce confusion in regards to this model selection. Although a discussion of the benefits and weaknesses of the multistage modeling approach is also included in the BMD guidance document, a summary of these should also be provided. The SAB also recommends that the draft assessment discuss the adequacy of the fit of the multistage model to available data. This discussion could be further

supported by exploring and reporting fits to other standard BMD model forms.

The SAB expresses concerns that the *assumption of independence of tumor incidence* that is required by the MS-COMBO methodology used in estimating the POD is not clearly described. In addition, there is no discussion on the validity of this assumption for the available RDX data. However, these data are available in the pathology report in Lish et al. (1984), and concurrent incidence of liver and lung tumors was found in only one animal in the 175/100 mg/kg-day dose group and in one animal in the 35 mg/kg-day dose group, demonstrating that the assumption of independence of tumor incidence holds. Finally, the SAB cannot determine whether the MS-COMBO methodology requires that the dose-response for each tumor be adequately described by a multistage model, and whether the tumor incidence data being combined must adequately fit the same multistage model form. The SAB recommends that a better, and more detailed description of the MS-COMBO methodology be provided in the draft assessment, and that this description clarify the points raised above. In particular, text that better describes the independence assumption and the impact of violations of this assumption on the estimated POD should be included.

### ***Key Recommendations***

- For liver cancer, perform and discuss results from additional BMD modeling (i.e. a sensitivity analysis documented in the Supplemental Materials) that examines and discusses:
  - The impact of low concurrent controls on model choice and the POD estimate.
  - The effect of including/excluding the highest dose has on model choice and the POD estimate.
- Provide details and discuss the adequacy of the fit of the multistage model to available data.
- Provide a better and more detailed description of the MS-COMBO methodology (in the Supplemental Materials) and ensure that this description discusses the issues below:
  - importance of the assumption of independence,
  - why this assumption is needed,
  - how this assumption might be examined statistically given adequate study data/documentation,
  - whether the independence of tumor incidence assumption further constrains the tumor-specific dose-response model form to be the same across included tumor types, and,
  - the extent to which violations of the independence of tumor incidence assumption negatively affect the estimated POD.

### ***Suggested Recommendations***

- Fit other standard BMD model forms to available data and include these findings as part of the discussion of model adequacy.
- Include a summary (in the Supplemental Materials) of the benefits and weaknesses of the multistage modeling approach.

### 3.4. Dose-Response Analysis

#### 3.4.1. Oral Reference Dose for Effects other than Cancer.

*Charge Question 4a. The draft assessment presents an overall oral reference dose of  $3 \times 10^{-3}$  mg/kg-day, based on nervous system effects as described in the Crouse et al. (2006) study. Is this selection scientifically supported and clearly described, including consideration of mortality as described in Section 2.1.6, and consideration of the organ/system-specific reference dose derived from the toxicity study by Cholakakis et al. (1980) that is lower (by approximately fivefold) as described in Section 2.1.4?*

EPA has done a reasonably good job describing the process and choices made to arrive at the oral RfD. The SAB concludes that the overall RfD for RDX should be based on nervous system effects. Neurotoxicity was observed in multiple animal studies and in exposed humans, and included hyperactivity, hyperirritability, tremors and convulsions. Mechanistic data supports the neurotoxic effects of RDX, namely binding to the GABA<sub>A</sub>R and antagonizing GABA-mediated post-synaptic inhibition. EPA also provides an RfD based on suppurative prostatitis and another based on testicular degeneration. The SAB finds that suppurative prostatitis, which EPA describes as a surrogate for the effects of RDX on the kidney and genitourinary system, is not an appropriate toxicological endpoint for the overall RfD. There is no known mechanistic link between suppurative prostatitis and renal papillary necrosis or adverse effects on renal function. Thus, suppurative prostatitis does not provide any indication of adverse effects in the kidneys. The SAB also concludes that testicular degeneration was not an appropriate endpoint to serve as a basis for the overall RfD. Testicular degeneration was reported at terminal sacrifice (i.e., 24 months) in one 2-year dietary study in mice (Lish et al. 1984). The SAB notes that testicular histopathology should have been seen at earlier time points (e.g., the six and 12 months' interim sacrifices) in Lish et al. (1984), as these exposure durations were several times longer than the 1.4-month duration of spermatogenesis in mice. Germ cell degeneration was also observed in a 2-year dietary study in rats (Levine et al. 1983) but only at the 12-month interim sacrifice, and not at the six-month interim or 24-month terminal sacrifice. Furthermore, testicular degeneration was not observed in most of the dietary and gavage studies in rodents (five of seven showed no effect).

Although the SAB agrees that neurotoxicity should be the basis for an overall RfD for RDX, the SAB concludes that the scientific support for the proposed oral RfD is weak.

EPA chose the Crouse et al. (2006) study over the Cholakakis et al. study for several stated reasons: lack of specific monitoring for neurological effects (e.g., convulsions) in the Cholakakis et al. study; a higher purity of test material in the Crouse study; fewer dose groups and wider spacing of dose groups in Cholakakis et al. (1980) compared to the Crouse study; and longer exposure duration in Crouse et al. (90 d) compared to Cholakakis et al. (14 d). While acknowledging that the Crouse study was a better designed study to detect neurological effects, and that the monitoring for neurological effects in Cholakakis et al. (1980) was incomplete, the observation of a (single) rat with convulsions at 2 mg/kg-day was a valid observation and the fact that it was noted despite incomplete observation suggests the possibility that more complete observation in this study could have identified an even lower LOAEL. The SAB acknowledges that the greater purity of the test material in the Crouse study is an important issue that should

1 impact the choice of the key study. The more informative dose spacing in Crouse et al. (2006)  
2 can, potentially, allow for less uncertainty in dose-response modeling. However, the SAB finds  
3 that the empirical observation of convulsion at the lower dose in Cholakakis et al. (1980) is also  
4 informative. The exposure duration was considerably longer in Crouse et al. (2006), and the  
5 potential advantage of longer exposure duration is that it may show effects at lower doses than in  
6 other studies of shorter duration even with the same dose range.

7  
8 EPA raised an additional quality consideration with respect to the Cholakakis et al. (1980)  
9 teratology study during the SAB review noting the observation of a single incident of convulsion  
10 in the positive control group treated with hydroxyurea. In the Cholakakis et al. study, RDX elicited  
11 convulsions in pregnant rats in a dose-related manner, consistent with other toxicological studies  
12 with RDX. Hydroxyurea, a known teratogen and consequently positive control substance, is also  
13 known to target the CNS (fetal and adult) (IARC, 2000). EPA cited the lack of convulsions in  
14 rats and dogs after repeated oral dosing in Morton et al. (2015), as evidence that hydroxyurea  
15 does not cause convulsions in laboratory animals. It should be noted, however, that other  
16 evidence of CNS stimulation; mainly aggression, was observed in this study in male rats given  
17 1,500 mg/kg-day hydroxyurea by gavage. Additionally, group sizes in the Morton et al. study  
18 were small (n= 3 to 5 per sex) such that the power of the study to identify a rare effect e.g.,  
19 convulsions, was insufficient. Hence, the Cholakakis et al. study observation of convulsions in one  
20 of the positive control animals is a plausible finding, and does not negate the convulsions  
21 observed in RDX treated animals.

22  
23 An additional study quality consideration regarding the Cholakakis et al. (1980) study is the  
24 potential lack of uniformity/homogeneity of the dosing preparations. In both the Cholakakis et al.  
25 (1980) teratology study and the Crouse et al. (2006) study, doses of RDX were administered in a  
26 suspension using methyl cellulose / Tween 80 as the vehicle. Cholakakis et al. acknowledges that  
27 “maintaining uniform suspensions was not always easy,” and when the same nominal  
28 concentration was assayed repeatedly, it showed wide variation in RDX content (33% to 500%  
29 relative to nominal). Less variability in RDX dose suspensions was reported by Crouse et al.  
30 (2006): “Each dose suspension was mixed using a magnetic stirring bar until a uniform  
31 suspension was obtained, and continued to be mixed each day during the dosing procedure.”  
32 Since these measures were taken by Crouse et al. to reduce the variation in dosing suspensions, it  
33 is likely that the intended dose levels were more accurately administered in the Crouse study  
34 compared with the Cholakakis study, where both under-dosing and over-dosing of animals is a  
35 concern due to the difficulty in maintaining uniform dose suspensions.

36  
37 Given the quality issues identified for the Cholakakis et al. (1980) study (with some of those issues  
38 articulated in the EPA draft report, and with SAB’s concern described above regarding the high  
39 variability of dose levels based on the difficulty in maintaining homogeneous dosing  
40 suspensions), it is appropriate to give more weight to the Crouse et al. (2006) study with respect  
41 to the quantitative dose-response analysis. However, the single incidence of convulsion observed  
42 in the 2 mg/kg-day RDX dose group in the Cholakakis et al. study should not be entirely  
43 discounted. Pregnant rats were used in the Cholakakis et al. study and males and non-pregnant  
44 female rats were used in the Crouse et al. study, but the available data do not resolve the question  
45 of whether pregnancy is a sensitive physiological state for the neurotoxicity of RDX. The

consideration that pregnancy may represent a sensitive physiological state should be considered in selecting the UF<sub>H</sub> for interspecies variability.

Nevertheless, the SAB considered options to specifically and quantitatively take the LOAEL from Cholakakis et al. (1980) into account.

1. Conduct a benchmark dose analysis on the convulsion data from Cholakakis et al. (1980).

EPA provided information that the incidence of convulsions at the high dose in the Cholakakis et al. study was combined with the incidence of other neurologic effects. The response at this dose is, therefore, not appropriate for inclusion in benchmark dose modeling. However, elimination of the high dose from the Cholakakis et al. dose-response data leaves only one effective dose and this does not provide an adequate basis for dose-response modeling. Therefore, the SAB rejects this option.

2. Combine the dose-response data from Cholakakis et al. (1980) and Crouse et al. (2006)

The SAB considered it may be possible to more specifically account for the data from pregnant dams in the Cholakakis et al. study by combining the data with those from Crouse et al. (2006). There were several impediments to this approach. The two studies differ in exposure duration (Cholakakis et al. (1980), 14-day; Crouse et al. (2006), 90-day) and sex/pregnancy status (Crouse et al. males and females; Cholakakis et al. pregnant females only) providing no common factors on which to combine results. Therefore, the SAB also rejects this option.

3. Use the NOAEL (0.2 mg/kg/d) from Cholakakis et al. (1980) as the POD

The SAB originally considered the advantages of using the NOAEL from Cholakakis et al. as a POD to derive the oral RfD, without full consideration of potential inaccuracies in the doses administered in the study. As noted above, with the elimination of the high dose from Cholakakis et al. (due to inclusion of non-convulsive effects), there is no basis for benchmark dose modeling, and a NOAEL is an appropriate basis for a POD. The use of the NOAEL from Cholakakis et al. as the basis for the RfD eliminates issues concerning the choice of a BMR from Crouse et al. (2006) (see response to charge question 3a(iii) in Section 3.3.1.3), and addresses the SAB's concern with the existence of a lower LOAEL from Cholakakis et al. However, given the quality issues associated with the Cholakakis et al. study, the SAB places more weight on the Crouse et al. (2006) study for the derivation of a POD, and therefore also rejects this option to use the NOAEL of the Cholakakis study.

4. Use the Dose-Response Data from Crouse et al. (2006) as the primary basis of deriving the RfD

The SAB recommends using the data from Crouse et al. (2006) in a benchmark dose analysis to derive a POD for the RfD. An RfD derived from the NOAEL of Cholakakis et al. (1980) can be used for comparison to the RfD derived from Crouse et al. Since the study population in Cholakakis et al. (1980) was a potentially sensitive subpopulation, an RfD of less than 10-fold lower derived from the NOAEL of Cholakakis et al. study is not unexpected.

Table 4 below provides a comparison between EPA's proposed value (first row entry) based on the 1% BMR from Crouse et al. (2006) with alternate RfDs. If the same UFs (composite UF of 100) are applied to the PBPK-adjusted NOAEL POD from the Cholakakis et al. study as EPA



applied to the PBPK-adjusted BMDL POD from the Crouse et al. study, the RfD based on Cholakis et al. (1980) would be  $1 \times 10^{-3}$  mg/kg/day. Applying the SAB recommended composite UF of 300 to the PBPK-adjusted NOAEL POD from Cholakis et al. (1980) results in an RfD of  $3 \times 10^{-4}$  mg/kg-day. Applying the SAB-recommended composite UF of 300 to the 1% BMR from Crouse et al. would result in an RfD of  $1 \times 10^{-3}$  mg/kg-day. If, however, the RfD from Crouse et al. were calculated based on a BMR of 5% as discussed by the SAB (see response to charge question 3a(iii)), applying the recommended composite UF of 300, the RfD would be  $4 \times 10^{-3}$  mg/kg-day. These RfDs can be compared to the EPA's proposed RfD of  $3 \times 10^{-3}$  mg/kg-day from Crouse et al. based on a BMR of 1% and a composite UF of 100.

**Table 4. Comparison of Derived Candidate RfD values using different PODs and composite uncertainty factors.**

Reference	POD (mg/kg-day)	POD Type	POD <sub>HED</sub> <sup>a</sup>	Composite UF	RfD value (mg/kg-day)
Crouse et al (2006)	0.57	BMDL <sub>01</sub>	0.28	100	0.003
Cholakis et al. (1980)	0.2	NOAEL	0.097	300	0.0003
Crouse et al. (2006)	0.569	BMDL <sub>01</sub>	0.28	300	0.001
Crouse et al. (2006)	2.66 <sup>b</sup>	BMDL <sub>05</sub>	1.295	300	0.004

<sup>a</sup> POD<sub>HED</sub> is calculated from POD x PBPK derived adjustment factor of 0.487

<sup>b</sup> BMDL<sub>05</sub> is from response to charge question 3a(iii)

#### Consideration of mortality

The SAB interprets this charge question as asking whether an RfD based on convulsions (from either Crouse et al. (2006), or Cholakis et al. (1980)) is adequately protective against lethality. The SAB agrees that mortality and convulsions are linked. However, the SAB is not aware of any evidence for RDX or similar seizurogenic compounds where neurologic mortality occurs in the absence of convulsions. The overall candidate RfDs (Table 4) can be compared to the NOAEL for convulsions of 10 mg/kg-day with no mortality in the monkey study of Martin and Hart (1974). The SAB finds that this comparison provides some confidence that an RfD based on a BMDL derived from Crouse et al. or the NOAEL from Cholakis et al. provides a margin of safety with respect to neurologic-based lethality. However, the SAB acknowledges that the Martin and Hart study had a small sample size. The SAB, therefore, strongly endorses increasing the UF<sub>D</sub> and apply a UF<sub>D</sub> of 10 to provide an appropriate margin of safety between convulsive and lethal neurologic effects (as well as accounting for data gaps in developmental neurotoxicity and lack of incidence data for less severe neurotoxic effects).

## Key Recommendations

- The SAB agrees that the overall RfD should be based on neurotoxicity, but the scientific support for the proposed oral RfD is weak.
- The discussion and key recommendations from Section 3.3.1.3 and Section 3.3.1.4 are all pertinent to the SAB finding that the scientific support for and discussion of the proposed oral RfD for the convulsions endpoint is weak. These recommendations are repeated here:
  - EPA should consider using a BMR of 5% for their dose-response modeling of the Crouse et al. (2006) data while addressing the uncertainty of using data on a frank effect (convulsions in this case) as the basis of an RfD with additional uncertainty factors.
  - If EPA decides to use a BMR of 1% for the dose-response assessment using Crouse et al. (2006), EPA should justify why the greater conservatism in risk assessment required for a frank effect (due to the lack of incidence data for less severe endpoints) is better dealt with through a lower BMR than through application of UF<sub>D</sub>.
  - If EPA decides to use a BMR of 1% for the Crouse et al. (2006), EPA should provide in its discussion clear justification for why a 1% BMR is more appropriate than a 5% BMR for RDX, given the greater uncertainty introduced into the dose-response assessment for RDX using a BMR of 1%.
  - Consistent with EPA guidance for UFs, the SAB strongly suggests that EPA apply a UF<sub>D</sub> of 10 to account for data gaps for developmental neurotoxicity, lack of incidence data for less severe neurological effects resulting in use of a severe effect (convulsions) as a basis for the RfD, and the proximity of lethal doses to convulsive doses.
  - EPA should discuss whether potential neurodevelopmental effects of RDX would be sufficiently addressed by a UF<sub>D</sub> of 10.
  - EPA should reconsider the value of the UFs and at a minimum provide stronger justification for a UF<sub>S</sub> of 1.
- Since the Cholakakis et al. (1980) study has several quality issues, it is appropriate to give more weight to the Crouse et al. (2006) study with respect to the quantitative dose-response analysis.

### 3.4.2. Inhalation Reference Concentration for Effects other than Cancer

*Charge Question 4b. The draft assessment does not derive an inhalation reference concentration as the available studies were insufficient to characterize inhalation hazard and conduct dose-response analysis, and no toxicokinetic studies of RDX were available to support development of a PBPK inhalation model. If you believe that the available data might support an inhalation reference concentration, please describe how one might be derived.*

There are no toxicokinetic data from inhalation exposure of laboratory animals or humans to RDX. There are epidemiological studies of persons exposed occupationally to RDX (Ma and Li, 1993; Hathaway and Buck, 1977), but no information is provided on exposure levels. These workers were likely exposed dermally and by inhalation.

**Key Recommendation:**

- EPA should not attempt to derive an inhalation reference concentration since neither toxicokinetic data nor exposure levels information from animal or human RDX inhalation studies are available to make estimation possible.

**3.4.3.Oral Slope Factor for Cancer**

*Charge question 4c. The draft assessment presents an overall oral slope factor of 0.038 per mg/kg-day based on the combination of liver and lung tumors in female mice. Is this derivation scientifically supported and clearly described?*

The SAB finds that the calculation of an oral slope factor (OSF) for cancer endpoints in the draft assessment is not clearly described and, in keeping with the discussion in question 3e(iii) (see Section 3.3.5.3), the SAB expresses several concerns below regarding whether the OSF is scientifically supported. The SAB makes multiple suggestions for how the discussion on the derivation of the OSF can be improved.

The OSF is estimated as the plausible upper-bound (95% upper CI) for the true slope, or risk per unit dose, from which the probability that an individual will develop cancer if exposed to an agent for a lifetime of 70 years can be derived. In practice, and as presented in the draft assessment, the OSF for the cancer endpoint is obtained as the slope of the line from a point of departure, in this case the BMDL for 10% BMR (BMDL<sub>10</sub>), to the estimated control response at a dose of zero. Consequently, any changes to the derivation of the POD will be reflected in the estimate of the OSF. In its response to question 3e(iii), the SAB identifies issues with the data used to derive the cancer POD, and offers recommendations for improving the calculation of the POD (see Section 3.3.5.3). These recommendations will change the estimated POD and thus the OSF.

The draft assessment proposes combining tumors from different sites in determining an overall cancer risk. The SAB finds that this is both logically and toxicologically sound. While not discussed in either the draft assessment or the draft supplemental material, the independence of tumor location is a key assumption for scientifically appropriate use of the MS-COMBO model. The SAB considered the original study data provided in the draft assessment, and agrees with the agency that there is no biological or statistical support for the notion that the two tumors used in the MS-COMBO analysis are interdependent. Hence the agency's assumption of independence of the two tumor sites and the MS-COMBO approach are considered valid. However, as discussed in the SAB's response to question 3e(iii), some issues around the use of the MS-COMBO model remain to be clarified.

The SAB expresses concern that the near linearity of the fitted multi-stage dose-response models (see Table 2-7 in the draft assessment identifying all selected models as Multistage 1°) results in a relatively poor fit (model estimates) at the highest doses. The two fitted models used (see Figures D-12 and D-14 in the RDX draft supplement document) have BMDL<sub>10</sub> estimates that are larger than the two lowest non-zero doses used. The Cancer Guidelines (USEPA, 2005) states (page 3-16): "If the POD is above some data points, it can fail to reflect the shape of the dose-response curve at the lowest doses and can introduce bias into subsequent extrapolations." This

seems to be what is happening with the RDX-induced adenomas and carcinomas data, and the issues with the BMDL<sub>10</sub> seem to arise primarily because the fitted multistage models (with parameter constraints invoked) lack sufficient curvature. Larger than expected BMDL<sub>10</sub> values (the PODs) result in lower estimated OSFs. The SAB conjectures that using a model form that allows more curvature could provide a better fit at the mid-range and higher doses, and improve the quality of fit. As mentioned in the response to question 3e(iii), the SAB acknowledges that EPA standard practice is to use the multistage model for benchmark dose modeling of cancer dose-response when there is no biological basis for choosing another model. In this case however, the relatively poor fit of the multistage model to the hepatocellular and alveolar/bronchiolar adenomas and carcinomas data produces an estimate of the POD with poor properties. The SAB recommends that at a minimum the draft assessment discuss the adequacy of the fit of the multistage model to available data. This discussion could be further supported by exploring and reporting fits to other standard BMD model forms – engaging in a curve-fitting exercise starting for example with the list in Table D-13 in the draft supplemental document. Although the multi-stage model does ensure positive slopes throughout, the BMDS software facilitates fitting other models that also adhere to this constraint.

The SAB expresses concern that the results for liver cancer in concurrent female control mice were very low (1.5 %) compared to available historical control incidence (8%; range 0-20%) (page 1-62 of the draft assessment). This low rate influences the final model for liver tumor incidence, which in turn significantly impacts the estimate of the POD and hence the OSF. It is not clear how this issue impacts the POD estimate derived via the MS-COMBO analysis where liver tumor results are combined with those of lung tumors to produce the final POD used. The SAB recommends that EPA acknowledge the low concurrent control liver tumor incidence in female mice and discuss its impact on the level of confidence in the final MS-COMBO estimate of the proposed POD.

The SAB also notes, and the draft assessment confirms (Section 1.2.5, page 1-61), that at the highest dose level in the Lish et al. (1984) study for the first 11 weeks, the animals had an elevated mortality strongly suggesting that the maximum tolerated dose had been greatly exceeded. At 11 weeks, the researchers lowered the dose, and it was a duration-weighted average dose level that was used as the highest dose in the fitting of the multistage model (see Section D.2.2 (pages D-31 to D-33) of the RDX draft supplement document). The Cancer Guidelines (page A-4) discuss this situation and offer that the decision to use or not use data from doses that exceed the MTD is “a matter of expert judgement”. The SAB has concerns that including this dose significantly impacts the final estimated POD. The SAB recommends that additional insight be sought by fitting the multistage model and estimating the POD after exclusion of this dose level, comparing the POD generated from both models, and discussing why the estimate that is based on the use of the highest dose data is preferred. Following this analysis through to the MS-COMBO results seems scientifically appropriate. This comparison and subsequent discussion is supported by the fact that the current IRIS entry for RDX of 0.11 per mg/kg-day was determined using the liver tumor data (Lish et al. 1984) with the highest dose values excluded.

#### **Key Recommendations**

- Acknowledge that the issues with the estimation of the POD for estimation of the cancer slope factor discussed in Section 3.3.5.3 and associated key recommendations for improving the presentation on the POD also apply to the estimation of the cancer slope factor.

#### **3.4.4. Inhalation Unit Risk for Cancer**

*Charge Question 4d. The draft assessment does not derive an inhalation unit risk because inhalation carcinogenicity data were not available, nor were toxicokinetic studies of inhalation of RDX available to support development of an inhalation PBPK model. If you believe that the available data might support an inhalation unit risk, please describe how one might be derived.*

There are no toxicokinetic data from inhalation studies of RDX in laboratory animals or humans, no inhalation carcinogenicity bioassays of RDX, nor data on cancer incidence in humans. Therefore, an inhalation unit risk for cancer cannot be derived.

#### **Key Recommendations**

- EPA should not attempt to derive an inhalation unit risk since there are no study data available to make estimation possible.

#### **3.5. Executive Summary**

*Charge Question 5. Does the executive summary clearly and adequately present the major conclusions of the assessment?*

Generally, the SAB considered the Executive Summary to be well-written, succinct, and clear. As changes are made to the body of the draft assessment in response to the SAB's recommendations, the Executive Summary should be updated accordingly. In addition, the SAB provides the following suggested recommendations for improving the Executive Summary.

#### ***Suggested Recommendations:***

- On the characterization and description of urogenital system hazard and risk.
  - Reduce the emphasis on the incidence and significance of the suppurative prostatitis. The other urogenital system effects are of more importance and should be described. The description of the urogenital effects in male rats should include specific mention of the renal effects (i.e. renal papillary necrosis and associated renal inflammation), not only the prostatic effects.
  - An RfD based on suppurative prostatitis should be derived as a stand-alone endpoint, and a separate RfD should be derived for kidney and other urogenital system effects. Because the observed suppurative prostatitis is part of a larger spectrum of prostatic inflammatory changes that are frequently found in aged F344 rats, the dose-response for this lesion found in male rats may in fact not reflect the overall incidence of all types of prostatitis combined in each dose group. The prostatic inflammation and renal/bladder effects may be inter-related, but this only occurs at the highest RDX dose tested, and there seems to be no basis for the assertion that suppurative prostatitis is a "surrogate marker" for renal/bladder effects. This change does not affect the overall oral reference dose because that is based on the nervous system effects.

- In the section on “Suppurative prostatitis,” the possibility of a bacterial infection is raised and its potential significance to RDX toxicity is briefly discussed. However, it should also be noted that this inflammatory effect could also occur without a bacterial infection.
  - P xxiii line 7 – 9: The first sentence indicates human potential for kidney and urogenital toxicity, which is justified, but indicates this is “based on” increased relative kidney weights and histopathological changes. P 1-24 lines 24-30 indicates inconsistent findings in the subchronic studies and down-plays the organ weight findings in the chronic studies, so the executive summary is inconsistent with this.
- Regarding the description of animal cancer bioassay results, the following should be added to indicate some of the uncertainty or limitations in the animal cancer bioassay results.
    - In the Summary, add “limited” to the sentence -. *“Results from animal studies provide suggestive evidence of carcinogenic potential for RDX based on **limited** evidence of positive trends in liver and lung tumor incidence in experimental animals.”*
    - In the body, add clarifying or cautionary language on page xxv, line 26 such as *Despite limitations in the animal cancer studies, a quantitative estimate of carcinogenic risk....* or *Cognizant of limitations in the animal cancer studies, a quantitative estimate of carcinogenic risk....*
  - On other content clarification, the following missing information should be included and the following editorial comments should be addressed:
    - P xxvii line 23 – 25: Please clarify the meaning of “more representative of potential human exposures,” and be explicit regarding the uncertainty associated with identifying a representative experimental exposure. It is not clear, given the limited information on RDX exposures in the Preface or elsewhere in the draft assessment, that dietary exposure is “*more representative of potential human exposures*”. It seems possible that human exposures could involve different or varied sources of RDX exposure (e.g., on swallowed dust particles, in consumed soil, incorporated into plants) such that neither experimental exposures as diet nor as gavage would be obviously “representative” and both experimental approaches to exposure (dosing) would include uncertainty.
    - Explain the importance of *RDX purity* in published studies.
    - List the main criteria used in choosing the principal study.
    - Include a discussion of the concordance in doses producing convulsions and doses at which death occurred in key animal studies in the brief discussion of neurologic effects in the section entitled “Effects other than cancer observed during oral exposure.” The lethality associated with convulsions is currently not mentioned.
    - Provide a summary statement addressing the confidence (i.e., low, medium or high) in the RfD.
    - On page xxiii, a paragraph break is needed after the sentence, “*There is no known MOA for male reproductive effects of RDX exposure.*” The next sentence does not relate to the male reproductive effects but speaks to the evidence for effects in other organs/systems.
    - Combine the paragraphs found on page xxv entitled “Effects other than cancer observed following inhalation exposure” and “Inhalation reference concentration (RfC) for effects other than cancer.” There is no available literature to support the identification of hazards following inhalation and a reference concentration cannot be determined. This can be stated simply in a single paragraph.

## REFERENCES

- Angerhofer, R; Davis, G; Balezewski, L. (1986). Teratological assessment of Trinitro-RDX in rats. (75-5100573-86). Aberdeen Proving Ground, MD: U.S. Army Environmental Hygiene Agency.
- Arfsten, DP; Davenport, R; Schaeffer, DJ. (1994). Reversion of bioluminescent bacteria (Mutatox) to their luminescent state upon exposure to organic compounds, munitions, and metal salts. *Biomed Environ Sci* 7, 144-149.
- Bach, PH; Nguyen, TK. (1998) Renal papillary necrosis--40 years on. *Toxicol Pathol.* 26(1):73-91.
- Bannon, DI; Dillman, JF; Phillips, CS; Perkins, EL. (2009). Global gene expression in rat brain and liver after oral exposure to the explosive hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX). *Chem Res Toxicol* 22: 620-625.
- Beyer, BK; Chernoff, N; Danielsson, BR; Davis-Bruno, K; Harrouk, W; Hood, RD; Janer, G; Wandel LU; Kim, JH; Rocca M; Rogers J; Scialli, AR. (2011) ILSI/HESI Maternal Toxicity Workshop Summary: Maternal Toxicity and Its Impact on Study Design and Data Interpretation. *Birth Defects Research (Part B)*. 92:36–51.
- Bhushan, B; Trott, S; Spain, JC; Halasz, A; Paquet, L; Hawari, I. (2003). Biotransformation of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) by a rabbit liver cytochrome P450: Insight into the mechanism of RDX biodegradation by Rhodococcus sp. Strain DN22. *Appl Environ Microbio* 69: 1347-1351. <http://dx.doi.org/10.1128/AEM.69.3.1347-1351.2003>
- Brix, AE. (2002) Renal papillary necrosis. *Toxicol Pathol.*30:672-674.
- Bronson, FH; Desjardins, C. (1977). Reproductive failure in aged CBF<sub>1</sub> male mice: Interrelationships between pituitary gonadotropic hormones, testicular function, and mating success. *Endocrinology* 101: 939-945.
- Bruni, JF; Huang, HH; Marshall, S; Meites, J. (1977). Effects of single and multiple injections of synthetic GnRH on Serum LH, FSH and testosterone in young and old male rats. *Biol Reprod* 17: 309-312.
- Bruyns-Haylett, M; Luo, J; Kennerley, A; Harris, S; Boorman, L; Milne, E; Vautrelle, N; Hayashi, Y; Whalley, BJ; Jones, M; Berwick, J; Riera, J; Zheng, Y. (2017). The neurogenesis of P1 and N1: A concurrent EEG/LFP study. *NeuroImage* 146: 575-588.
- Cao, CI; Reddy, G; Bannon, DI; Hohnson, MS. (2008). In vitro study of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) metabolism in human liver. Aberdeen Proving Ground, ND: U.S. Army Center for Health Promotion and Preventive Medicine.

- 1 Carney, EW; Kimmel, CA, (2007) Interpretation of Skeletal Variations for Human Risk  
2 Assessment: Delayed Ossification and Wavy Ribs. *Birth Defects Research (Part B)-*  
3 *Developmental and Reproductive Toxicology* 80:473-496.  
4
- 5 Chen, H; Hardy, MP; Zirkin BR. (2002). Age-related decreases in Leydig cell testosterone  
6 production are not restored by exposure to LH in vitro. *Endocrinology* 143: 1637–1642.  
7
- 8 Cholakakis, JM; Wong, LCK; Van Goethem, DL; Minor, J; Short, R; Sprinz, H; Ellis, HV, III.  
9 (1980). Mammalian toxicological evaluation of RDX. (DAMD17-78-8027). Kansas City,  
10 MO: Midwest Research Institute.  
11
- 12 Cohen, BJ; Anver, MR; Ringler, DH; Adelman, RC. (1978). Age-associated pathological  
13 changes in male rats. *Federation Proc* 37: 2848-2850.  
14
- 15 Coleman, NV, Spain, JC, Duxbury, T. (2002). Evidence that RDX biodegradation by  
16 *Rhodococcus* strain DN22 is plasmid-borne and involves a cytochrome p-450. *J Appl*  
17 *Microbiol* 93: 463-472.  
18
- 19 Costa, LG; Giordano, G; Faustman, EM. (2010). Domoic acid as a developmental neurotoxin.  
20 *Neurotoxicology* 33: 409-423.  
21
- 22 Creasy, D; Bube , A; de Rij, KE; Kandori, H; Kuwahara, M; Masson, R; Nolte, T; Reams, R;  
23 Regan, K; Rehm, S; Rogerson, P; Whitney, K. (2012). Proliferative and nonproliferative  
24 lesions of the rat and mouse male reproductive system. *Toxicol Pathol.* 40(6 Suppl):40S-  
25 121S.  
26
- 27 Creeley, CE. (2016). From drug-induced developmental neural apoptosis to pediatric anesthetic  
28 neurotoxicity – where are we now? *Brain Sciences* 6:32-44  
29
- 30 Crouse, LCB; Michie, MW; Major, M; Johnson, MS; Lee, RB; Paulus, HI. (2006). Subchronic  
31 oral toxicity of RDX in rats. (Toxicology Study No. 85-XC-5131-03). Aberdeen Proving  
32 Ground, MD: U.S. Army Center for Health Promotion and Preventive Medicine.  
33
- 34 Dey, A; Kang, X; Qiu, J; Du, Y; Jiang, J (2016). Anti-Inflammatory Small Molecules To Treat  
35 Seizures and Epilepsy: From bench to Bedside. *Trends in Pharmacological Sciences* 37:  
36 463-484.  
37
- 38 Doucette, TA; Tasker, R. (2016) Perinatal domoic acid as a neuroteratogen. *Curr Top Behav*  
39 *Neurosci.* 29:87-110.  
40
- 41 Eyo, U. B; Murugan, M; Wu, LJ. (2017), Microglia–Neuron communication in epilepsy. *Glia*  
42 65: 5–18. doi:10.1002/glia.23006  
43
- 44 Fellows, RJ; Driver, CR; Cataldo, DA; Harvey, SD. (2006). Bioavailability of hexahydro-1,3,5-  
45 trinitro-1,3,5-triazine (RDX) to the prairie vole (*Microtus ochrogaster*). *Environmental*  
46 *Toxicology and Chemistry* 25:1881-1886.



- 1
- 2 Fuller, ME; Perreault, N; Hawari, J. (2010). Microaerophilic degradation of hexahydro-1,3,5-
- 3 trinitro-1,3,5-triazine (RDX) by three Rhodococcus strains. *Letters in Applied*
- 4 *Microbiology* 51:313–318.
- 5
- 6 Fuller, ME; McClay, K; Hawari, J; Paquet, L; Malone, TE; Fox, BG; Steffan, RJ. (2009).
- 7 Transformation of RDX and other energetic compounds by xenobiotic reductases XenA
- 8 and XenB. *Appl Microbiol Biotechnol* 84:535-544.
- 9
- 10 Galineau, L; Kas, A; Worbe, Y; Chaigneau, M; Herard, AS; Guillermier, M; Delzescaux, T;
- 11 Féger J, Hantraye, P; Tremblay, L. (2017). Cortical areas involved in behavioral
- 12 expression of external pallidum dysfunctions: A PET imaging study in non-human
- 13 primates. *Neuroimage*.146:1025-1037.
- 14
- 15 Garcia-Reyero, N; Habib, T; Pirooznia, M; Gust, KA; Gong P; Warner, C; Wilbanks, M;
- 16 Perkins, E. (2011). Conserved toxic responses across divergent phylogenetic lineages: a
- 17 meta-analysis of the neurotoxic effects of RDX among multiple species using
- 18 toxicogenomics. *Ecotoxicology* 20:580-594.
- 19
- 20 George, SE; Huggins-Clark, G; Brooks, LR. (2001). Use of a Salmonella microsuspension
- 21 bioassay to detect the mutagenicity of munitions compounds at low concentrations. *Mutat*
- 22 *Res* 490: 45-56.
- 23
- 24 Gill, DA; Bastlund, JF; Watson, WP; Ryan, CL; Reynolds, DS; Tasker, RA. (2010). Neonatal
- 25 exposure to low-dose domoic acid lowers seizure threshold in adult rats. *Neuroscience* 169:
- 26 1789-1799.
- 27
- 28 Gosden, RG; Richardson, DW; Brown, N; Davidson, DW (1982). Structure and gametogenic
- 29 potential of seminiferous tubules in ageing mice. *J Reprod Fert* 64: 127-130.
- 30
- 31 Grant, KS; Burbacher, TM; Faustman, EM; Gratttan, L. (2010). Domoic acid: neurobehavioral
- 32 consequences of exposure to a prevalent marine biotoxin. *Neurotoxicol Teratol.* 32:132-
- 33 141.
- 34
- 35 Grasso, C; Li Volsi, G; Cataldo, E; Manzoni, D; Barresi, M. (2016). Effects
- 36 of bicuculline application on the somatosensory responses of secondary vestibular
- 37 neurons. *Neuroscience* 335:122-133.
- 38
- 39 Guo, L; Xu, H; Chen, Y; Chang, Y. (1985). Distribution and metabolism of tritium-labeled
- 40 hexogen in white mice, *Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Za Zhi* 3: 335.330.
- 41
- 42 Gust, KA; Brasfield, SM; Stanley, JK; Wilbanks, MS; Chappell, P; Perkins, EJ; Lotufo, GR;
- 43 Lance, RF. (2011). Genomic investigation of year-long and multigenerational exposures
- 44 of fathead minnow to the munition compound RDX. *Environmental Toxicology and*
- 45 *Chemistry* 30:1852-1864.
- 46

- 1 Halasz, A; Manno, D; Perreault, NN; Sabbadin, F; Bruce, NC; Hawari, J. (2012). Biodegradation  
2 of RDX Nitroso Products MNX and TNX by Cytochrome P450 XplA. *Environmental*  
3 *Science and Technology* 46:7245-7251.  
4
- 5 Hart, ER. (1976). Two-year chronic toxicity study in rats. (N0014-73-C-0162). Kensington, MD:  
6 Litton Bionetics, Inc.  
7
- 8 Haseman, JK; Huff, JE; Rao, GN; Arnold, JE; Boorman, GA; McConnell, EE. (1985).  
9 Neoplasms observed in untreated and corn oil gavage control groups of F344/N rats and  
10 (C57BL/6N X C3H/HeN)F1 (B6C3F1) mice. *J Natl Cancer Inst* 75: 975-984.  
11
- 12 Hathway, JA; Buck, CR. (1977). Absence of health hazards associated with RDX manufacture  
13 and use. *J Occup Med* 19: 269-272.  
14
- 15 Hess-Ruth, A; Crouse, L; Roszell, L. (2007). RDX pilot development neurotoxicity test in rats.  
16 (Toxicology Study No. 85-XC-064Y-07). Aberdeen Proving Grounds: U.S. Army Center  
17 for Health Promotion and Preventive Medicine.  
18
- 19 Hiolski, EM; Ito, S; Beggs, JM; Lefebvre, KA; Litke, AM; Smith, DR. (2016). Domoic  
20 acid disrupts the activity and connectivity of neuronal networks in organotypic brain slice  
21 cultures. *Neurotoxicology* 56:215-224.  
22
- 23 Hollander, AI; Colbach, EM. (1969). Composition C-4 induced seizures: A report of five cases.  
24 *Mil Med* 134: 1529-1530.  
25
- 26 Hu, Z; Li, Z (2017). miRNAs in synapse development and synaptic plasticity. *Curr Opin*  
27 *Neurobiol* 45: 24-31.  
28
- 29 IARC (International Agency for Research on Cancer). (2000). Some antiviral and antineoplastic  
30 drugs, and other pharmaceutical agents. *IARC Monographs on the evaluation of*  
31 *carcinogenic risks to humans* Vol 76. Pg 347 - 386.  
32 <https://monographs.iarc.fr/ENG/Monographs/vol76/mono76-14.pdf>;  
33
- 34 Jaligama, S; Kale, VM; Wilbanks, MS; Perkins, EJ; Meyer, SA. (2013). Delayed  
35 myelosuppression with acute exposure to hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX)  
36 and environmental degradation product hexahydro-1-nitroso-3,5-dinitro-1,3,5-triazine  
37 (MNX) in rats. *Toxicology and Applied Pharmacology* 266:443-451.  
38
- 39 Jeilani, YA; Duncan, KA; Newallo, DS; Thompson, AN Jr.; Bose, NK. (2015). Tandem mass  
40 spectrometry and density functional theory of RDX fragmentation pathways: Role of ion-  
41 molecule complexes in loss of NO<sub>3</sub> and lack of molecular ion peak. *Rapid*  
42 *Communications in Mass Spectrometry* 29:802-810.  
43
- 44 Johnson, L; Neaves, WB. (1983). Enhanced daily sperm production in the remaining testis of  
45 aged rats following hemicastration. *J Androl* 4:162-166.

- Kim, JY; Liu, CY; Zhang, F; Duan, X; Wen, Z; Song, J; Feighery, E; Lu, B; Rujescu, D; St Clair D; Christian, K; Callicot, JH; Weinberger, DR; Song, H; Ming, GL. (2012). Interplay between DISC1 and GABA signaling regulates neurogenesis in mice and risk for schizophrenia. *Cell* 148:1051-1064.
- Krishnan, K; Crouse, LCB; Bazar, MA; Major, MA; Reddy, G. (2009). Physiologically based pharmacokinetic modeling of cyclotrimethylenetrinitramine in male rats. *J Appl Toxicol* 29: 629-637. <http://dx.doi.org/10.1002/jat.1455>
- Kucukardali, Y; Acar, HV; Ozkan, S; Nalbant, S; Yazgan, Y; Atasoy, EM; Keskin, O; Naz, A; Akyatan, N; Gokben, M; Danaci, M. (2003). Accidental oral poisoning caused by RDX (cyclonite): A report of 5 cases. *J Intensive Care Med* 18: 42-46. <http://dx.doi.org/10.1177/0885066602239123>
- Lee, E; Lee, J; Kim, E. (2016) Excitation/Inhibition Imbalance in Animal Models of Autism Spectrum Disorders. *Biol Psychiatry* 81: 838-847.
- Levine, BS; Furedi, EM; Gordon, DE; Burns, JM; Lish, PM. (1981a). Thirteen week toxicity study of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) in Fischer 344 rats. *Toxicol Lett* 8: 241-245.
- Levine, BS; Furedi, EM; Gordon, DE; Burns, JM; Lish, PM. (1981b). Thirteen week oral (diet) toxicity study of trinitrotoluene (TNT) and RDX in F344 rats. *Fundam Appl Toxicol* 15: 373-380. [http://dx.doi.org/10.1016/0272-0590\(90\)90062-0](http://dx.doi.org/10.1016/0272-0590(90)90062-0)
- Levine, BS; Lish, PM; Furedi, EM; Rac, VS; Sagartz, JM. (1983). Determination of the chronic mammalian toxicological effects of RDX (twenty-four month chronic toxicity/carcinogenicity study of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) in the Fischer 344 rat): Final report—phase V, Chicago, IL: IIT Research Institute.
- Levine, BS; Furedi, EM; Gordon, DE; Barkley, JJ; Lish, PM. (1990). Toxic interactions of the munitions compounds TNT and RDX in F344 rats. *Fundam Appl Toxicol* 15: 373-380.
- Lish, PM; Levine, BS; Furedi, EM; Sagartz, JM; Rac, VS. (1984) Determination of the chronic mammalian toxicological effects of RDX: Twenty-four month chronic toxicity/carcinogenicity study of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) in the B6C3F1 hybrid mouse (Volumes 1 and 3). U.S. Army Medical Research and Development Command, Chicago, IL.
- Ma, B; Li, H. (1993). Neurobehavior effects of hexogen on exposed workers. *Gongye Weisheng yu Zhiyejin* 19: 20-23.
- Major, MA; Reddy, G; Berge, MA; Patzer, SS; Li, AC; Gohdes, M. (2007). Metabolites profiling of [14C] hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) in Yucatan miniature pigs. *J Toxicol Environ Health A* 70: 1191-1202. <http://dx.doi.org/10.1080/15287390701252717>

- 1 Marriott, AL; Tasker, RA; Ryan, CL; Doucette, TA. (2016). Alterations to prepulse inhibition  
2 magnitude and latency in adult rats following neonatal treatment with domoic acid and  
3 social isolation rearing. *Behav Brain Res* 298(Pt B):310-317.  
4
- 5 Martin, D; Hart, E. (1974). Subacute toxicity of RDX and TNT in monkeys (pp. 1-216).  
6 (ADA044650). Kensington, MD: Litton Bionetics, Inc.  
7
- 8 Marty, S; Wehrle, R; Sotelo, C. (2000). Neuronal activity and brain-derived neurotrophic factor  
9 regulate the density of inhibitory synapses in organotypic slice cultures of postnatal  
10 hippocampus. *The Journal of Neuroscience* 20:8087-8095.  
11
- 12 McCormick, NG; Cornell, JH; Kaplan, AM. (1981). Biodegradation of hexahydro-1,3,5-trinitro-  
13 1,3,5-triazine. *Appl Environ Microbiol* 42: 817-823.  
14
- 15 Melnick, RL; Thaye, KA; Buche, JR. (2008). Conflicting views on chemical carcinogenesis  
16 arising from the design and evaluation of rodent carcinogenicity studies. *Environ Health*  
17 *Perspect* 116: 130-135.  
18
- 19 Merrill, SL. (1968). Ingestion of an explosive material, composition C-4: A report of two cases.  
20 *USARV Med Bull* 40: 5-11.  
21
- 22 Meunier, CN; Chameau, P; Fossier, PM. (2017). Modulation of Synaptic Plasticity in the Cortex  
23 Needs to Understand All the Players. *Front Synaptic Neurosci* 9:2-15. doi:  
24 10.3389/fnsyn.2017.00002.  
25
- 26 Meyer, SA; Marchand AJ; Hight JL; Roberts GH; Escalon LB; Inouye LS; MacMillan DK. (2005).  
27 Up-and-down procedure (UDP) determinations of acute oral toxicity of nitroso degradation  
28 products of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX). *Journal of Applied Toxicology*  
29 25:427-434.  
30
- 31 Mills, BD; Pearce, HL; Khan, O; Jarrett, BR; Fair, DA; Lahvis, GP. (2016). Prenatal domoic  
32 acid exposure disrupts mouse pro-social behavior and functional connectivity MRI.  
33 *Behav Brain Res* 308:14-23.  
34
- 35 Morton, D; Reed, L; Huang, W; Marcek, JM; Austin-LaFrance, R; Norithcott, CA; Schelling,  
36 SH; Enerson, BE; Tomlinson, L. (2015). Toxicity of hydroxyurea in rats and dogs.  
37 *Toxicologic Pathology* 45: 498-512.  
38
- 39 Mukhi, S; Patino, R. (2008). Effects of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) in  
40 zebrafish: General and reproductive toxicity. *Chemosphere* 72:726-732.  
41
- 42 Musick, TI; Berge, MA; Patzer, SS; Tilch, KR. (2010). Absorption, distribution, metabolism,  
43 and excretion of <sup>14</sup>C-RDX following oral administration to minipigs (DAAD05-02-P-  
44 2319). (ADA526472). Madison, WI: Covance Laboratories Inc. [http://www.dtic.mil/cgi-](http://www.dtic.mil/cgi-bin/GetTRDoc?AD=ADA526472)  
45 [bin/GetTRDoc?AD=ADA526472](http://www.dtic.mil/cgi-bin/GetTRDoc?AD=ADA526472)  
46

- 1 Nasrallah, FA; Kaur, K; Yeow, LY; Chuang, KH. (2017). GABAergic effect on resting-state  
2 functional connectivity: Dynamics under pharmacological antagonism. *NeuroImage* 149:  
3 53-62.  
4
- 5 Nasehi, M; Roghani, F; Ebrahimi-Ghiri, M; Zarrindast, MR. (2017). Role of the amygdala  
6 GABA-A receptors in ACPA-induced deficits during conditioned fear learning. *Brain*  
7 *Res Bull* 131:85-92.  
8
- 9 Neal, GE; Judah, DJ; Hard, GG; Ito, N. (2003). Differences in ethoxyquin nephrotoxicity  
10 between male and female F344 rats. *Food Chem Toxicol* 41:193-200.  
11
- 12 Neal, AP; Guilarte, TR. (2013). Mechanism of lead and manganese neurotoxicity. *Toxicol Res*  
13 2:99-114.  
14
- 15 OECD (Organization for Economic Cooperation and Development) (2015). OECD guideline for  
16 testing of chemicals: Reproduction/Developmental Toxicity Screening Test (Test  
17 Guideline 421).  
18
- 19 Pan, X; M.J. San Francisco, MI; Lee, C; Ochoa, KM; Xu, X; Liu, J; Zhang, B; Cox, SB; Cobb,  
20 GP. (2007a) Examination of the mutagenicity of RDX and its N-nitroso metabolites using  
21 the Salmonella reverse mutation assay. *Mutat Res* 629, 64-69.  
22
- 23 Pan, X; Zhang, B; Smith, JN; San Francisco, M; Anderson, TA; Cobb, GP. (2007b). N-Nitroso  
24 compounds produced in deer mouse (*Peromyscus maniculatus*) GI tracts following  
25 hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) exposure. *Chemosphere* 67: 1164-1170.  
26 <http://dx.doi.org/10.1016/j.chemosphere.2006.10.077>  
27
- 28 Pan X; Ochoa KM; San Francisco MJ; Cox SB; Dixon K; Anderson TA; Cobb GP. (2013).  
29 Absorption, distribution, and biotransformation of hexahydro-1,3,5-trinitro-1,3,5-triazine  
30 in B6C3F1 mice (*Mus musculus*). *Environmental Toxicology and Chemistry* 32:1295-  
31 1303.  
32
- 33 Parker, GA; Reddy, G; Major, MA. (2006). Reevaluation of a twenty-four-month chronic  
34 toxicity/carcinogenicity study of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) in the  
35 B6C3F1 hybrid mouse. *Int J Toxicol* 25: 373-378.  
36
- 37 Reddy, G; Erexson, GL; Cifone, MA; Major, MA; Leach, GL. (2005) Genotoxicity assessment  
38 of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX). *Int J Toxicol* 24: 427-434.  
39
- 40 Rivera, C; Voipio, J; Payne, JA; Ruusuvuori, E; Lahtinen, H; Lamsa, K; Pirvola, U; Saarma, M;  
41 Kaila, K. (1999). The K<sup>+</sup>/Cl<sup>-</sup> co-transporter KCC2 renders GABA hyperpolarizing  
42 during neuronal maturation. *Nature* 397:251-255.  
43
- 44 Rogers, JM; Chernoff, N; Keen, CL; Daston, GP. (2005). Evaluation and interpretation of  
45 maternal toxicity in Segment II studies: Issues, some answers, and data needs. *Toxicology*  
46 *and Applied Pharmacology* 207: S367-74.

- Salari, AA; Amani, M. (2017) Neonatal blockade of GABA-A receptors alters behavioral and physiological phenotypes in adult mice. *Int J Dev Neurosci* 57:62-71.
- Schneider, NR; Bradley, SL; Andersen, ME. (1978). Distribution and metabolism of cyclotrimethylenetrinitramine (RDX) in rat after sub-chronic administration. *Toxicol Appl Pharmacology* 46: 163-171.
- Smith, JN; Pan, XP; Gentles, A; Smith, EE; Cox, SB; Cobb, GE. (2006). Reproductive effects of hexahydro-1,3,5-trinitroso-1,3,5-triazine in deer mice (*Peromyscus maniculatus*) during a controlled exposure study. *Environ Toxicol Chem* 25: 446-451.
- Smith, JN; Espino, MA; Liu, J; Romero, NA;; Cox, SB; Cobb, GP. (2009). Multigenerational effects in deer mice (*Peromyscus maniculatus*) exposed to hexahydro-1,3,5-trinitroso-1,3,5-triazine (TNX). *Chemosphere* 75: 910-914.
- Snodgrass, HL, Jr. (1984). Preliminary assessment of relative toxicity and mutagenicity potential of 1-nitroso-3,5-dinitro-1,3,5-triazacyclohexane (mononitroso-RDX) (Final Report ed.). (ADA149351. USAEHA-75-51-0345-85). Aberdeen Proving Ground, MD: U.S. Army Environmental Hygiene Agency.  
<http://oai.dtic.mil/oai/oai?verb=getRecord&metadataPrefix=html&identifier=ADA149351>
- Stamou, M; Streifel, KM; Goines, PE; Lein, PJ. (2013). Neuronal connectivity as a convergent target of gene  $\times$  environment interactions that confer risk for Autism Spectrum Disorders. *Neurotoxicol Teratol* 36:3-16.
- Suzuki, N; Withers HR. (1978). Exponential decrease during aging and random lifetime of mouse spermatogonial stem cells. *Science* 202: 1214-1215.
- Suwa, T; Nyska, A; Peckham, JC; Hailey, JR; Mahler, JF; Haseman, JK; Maronpot, RR..(2001). A retrospective analysis of background lesions and tissue accountability for male accessory sex organs in Fischer-344 rats. *Toxicol Pathol.* 29:467-78. PubMed PMID: 11560252.
- Sweeney, LM; Gut, CP, Jr; Gargas, ML; Reddy, G; Williams, LR; Johnson, MS. (2012a). Assessing the non-cancer risk for RDX (hexahydro-1,3,5-trinitro-1,3,5-triazine) using physiologically based pharmacokinetic (PBPK) modeling [Review] *Regul Toxicol Pharmacol* 62: 107-114. <http://dx.doi.org/10.1016/j.yrtph.2011.12.007>
- Sweeney, LM; Okolica, MR; Gut, CP, Jr; Gargas, ML. (2012b). Cancer mode of action, weight of evidence, and proposed cancer reference value for hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX). *Regul Toxicol Pharmacol* 64: 205-224.

- 1 Takahashi, K; Maejima, H; Ikuta, G; Mani, H; Asaka, T. (2017). Exercise combined with low-  
2 level GABA<sub>A</sub> receptor inhibition up-regulates the expression of neurotrophins in the  
3 motor cortex. *Neuroscience Letters* 636:101-107.  
4
- 5 Tan, EL; Ho, CH; Griest, WH; Tyndall, RL. (1992). Mutagenicity of trinitrotoluene and its  
6 metabolites formed during composting. *J Toxicol Environ Health A* 36: 165-175.  
7 <http://dx.doi.org/10.1080/15287399209531632>  
8
- 9 Testud, F; Glanclaude, JM; Descotes, I. (1996). Acute hexogen poisoning after occupational  
10 exposure. *J Toxicol Clin Toxicol* 34: 109-111.  
11 <http://dx.doi.org/10/3109/15563659609020244>  
12
- 13 U.S.EPA (Environmental Protection Agency). (1998a). Health effects test guidelines: OPPTS  
14 870.3700. Prenatal Developmental Toxicity Study. Washington, DC, United States  
15 Environmental Protection Agency, Office of Prevention, Pesticides and Toxic Substances  
16 (EPA/712/C- 98/207;  
17
- 18 U.S.EPA (Environmental Protection Agency). (1998b). Health effects test guidelines: OPPTS  
19 870.7800. Immunotoxicity. Washington, DC, United States Environmental Protection  
20 Agency, Office of Prevention, Pesticides and Toxic Substances (EPA/712/C- 98/351;  
21 [http://www.epa.gov/ocspp/pubs/frs/publications/Test\\_Guidelines/series870.htm](http://www.epa.gov/ocspp/pubs/frs/publications/Test_Guidelines/series870.htm))  
22
- 23 U.S.EPA (Environmental Protection Agency). (2005). Guidelines for Carcinogen Risk  
24 Assessment. (EPA/630/P-03/0001F). Washington, DC U.S. Environmental Protection  
25 Agency, Risk Assessment Forum.  
26
- 27 U.S.EPA (Environmental Protection Agency). (2011). Recommended use of body weight<sup>3/4</sup> as  
28 the default method in derivation of the oral reference dose. (EPA/100/R11/0001).  
29 Washington, DC. U.S. Environmental Protection Agency, Risk Assessment Forum.  
30
- 31 U.S.EPA (2012a). Benchmark dose technical guidance. (EPA/100/R-12/001). Washington, DC.  
32 U.S. Environmental Protection Agency. Risk Assessment Forum.  
33
- 34 U.S. EPA (2012b). Peer review summary report. External letter peer review of study reports on  
35 1,3,5-trinitro-1,3,5-hexahydrotriazine (RDX). Washington, DC. U.S. Environmental  
36 Protection Agency.  
37
- 38 Von Oettingen, WF; Donahue, DD; Yagoda, H; Monaco, AR; Harris, MR. (1949). Toxicity and  
39 potential dangers of cyclotrimethylenetrinitramine (RDX). *J Ind Hyg Toxicol* 31: 21-31.  
40
- 41 Wang, C; Leung, A; Sinhia-Hikim, AP. (1993). Reproductive aging in the male Brown-Norway  
42 rat: A model for the human. *Endocrinology* 133: 2773-2781.  
43
- 44 Ward, JM; Goodman, DG; Squire, RA; Chu, KC; Linhart, MS (1978). Neoplastic and  
45 nonneoplastic lesions in aging (C57BL/6N × C3H/HeN)F<sub>1</sub> (B6C3F<sub>1</sub>) mice. *JNCI* 83;  
46 849-854,



- West, RR; Stafford, DA. (1997) Occupational exposures and haematological abnormalities among ordnance factory workers: A case control study. *Leuk Res* 21: 675-680.
- WHO (World Health Organization). (2012). Guidance for immunotoxicity risk assessment for chemicals. (Harmonization Project Document No. 10). Geneva, Switzerland.  
<http://www.inchem.org/documents/harmproj/harmproj/harmproj10.pdf>
- Williams, LR; Bannon, DI. (2009). Mechanism of RDX-induced seizures in rats (Toxicology Study No. 87-XE-OBT9-09). (ADA515072. USACHPPM-TSN-87-XE-OBT9-09). Aberdeen Proving Ground, MD: US Army Center for Health Promotion and Preventive Medicine, Health Effects Research Program.  
<http://oai.dtic.mil/oai/oai?verb=getRecords&metadataPrefix=html&identifier=ADA511072>
- Williams, LR; Wong, K; Stewart, A; Suciu C; Gaikwad, S; Wu, N; DiLeo, J; Grossman, L; Cachat, J; Hart, P; Kalueff, AV. (2012). Behavioral and physiological effects of RDX on adult zebrafish. *Comparative Biochemistry and Physiology C-Toxicology and Pharmacology* 155:33-38.
- Williams, LR; Aroniadou-Anderjaska, V; Qashu, F; Finne, H; Pidoplichko, V; Bannon, DI; Braga, MF.(2011). RDX binds to the GABA(A) receptor-convulsant site and blocks GABA(A) receptor-mediated currents in the amygdala: a mechanism for RDX-induced seizures. *Environ Health Perspect.* 119: 357-363.
- Wirbisky, SE; Weber, GJ; Lee, JW; Cannon, JR; Freeman, JL. (2014) Novel dose-dependent alterations in excitatory GABA during embryonic development associated with lead (Pb) neurotoxicity. *Toxicol Letters* 229: 1-8.
- Woody, RC; Kearns, GL; Brewster, MA; Turley, CP; Sharp, GB; Lake, RS. (1986). The neurotoxicity of cyclotriethylenetrinitramine (RDX) in a child: A clinical and pharmacokinetic evaluation. *Clin. Toxicol* 24: 305-319.
- Zhang, X; Ebata, KT; Robaire, B; Nagano, MC. (2006). Aging of male germ line stem cells in mice. *Biol Reprod* 74: 119–124.
- Zhang, B; Pan, X. (2009). RDX Induces Aberrant Expression of MicroRNAs in Mouse Brain and Liver. *Env Health Perspect* 117: 231-240.
- Zuloaga, DG; Lahvis, GP; Mills, B; Pearce, HL; Turner, J; Raber, J. (2016). Fetal domoic acid exposure affects lateral amygdala neurons, diminishes social investigation and alters sensory-motor gating. *Neurotoxicology* 53:132-140.



## APPENDIX A: EPA'S CHARGE QUESTIONS

### Charge to the Science Advisory Board for the IRIS Toxicological Review of Hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) September 2016 ( Updated November 2016<sup>1</sup>)

#### Introduction

The U.S. Environmental Protection Agency (EPA) is seeking a scientific peer review of a draft Toxicological Review of Hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) developed in support of the Agency's online database, the Integrated Risk Information System (IRIS). IRIS is prepared and maintained by EPA's National Center for Environmental Assessment (NCEA) within the Office of Research and Development (ORD).

IRIS is a human health assessment program that evaluates scientific information on effects that may result from exposure to specific chemicals in the environment. Through IRIS, EPA provides high quality science-based human health assessments to support the Agency's regulatory activities and decisions to protect public health. IRIS assessments contain information for chemicals that can be used to support hazard identification and dose-response assessment, two of the four steps in the human health risk assessment process. When supported by available data, IRIS provides health effects information and toxicity values for health effects (including cancer and effects other than cancer) resulting from chronic exposure. IRIS toxicity values may be combined with exposure information to characterize public health risks of chemicals; this risk characterization information can then be used to support risk management decisions.

An existing assessment for RDX includes a reference dose (RfD) posted on the IRIS database in 1988 and oral slope factor (OSF) and a cancer descriptor posted in 1990. The IRIS Program is conducting a reassessment of RDX. The draft Toxicological Review of RDX is based on a comprehensive review of the available scientific literature on the noncancer and cancer health effects in humans and experimental animals exposed to RDX. Additionally, appendices for chemical and physical properties, toxicokinetic information, summaries of toxicity studies, and other supporting materials are provided as *Supplemental Information* (see Appendices A to D) to the draft Toxicological Review.

The draft assessment was developed according to guidelines and technical reports published by EPA (see *Preamble*), and contains both qualitative and quantitative characterizations of the human health hazards for RDX, including a cancer descriptor of the chemical's human carcinogenic potential, a noncancer toxicity value for chronic oral exposure (RfD), and a cancer risk estimate for oral exposure.

## Charge questions on the draft Toxicological Review of RDX

1. **Literature search/study selection and evaluation.** The section on *Literature Search Strategy/ Study Selection and Evaluation* describes the process for identifying and selecting pertinent studies. Please comment on whether the literature search strategy, study selection considerations including exclusion criteria, and study evaluation considerations, are appropriate and clearly described. Please identify additional peer-reviewed studies that the assessment should consider.
2. **Toxicokinetic modeling.** In Appendix C, Section C.1.5, the draft assessment presents a summary, evaluation, and further development of published PBPK models for RDX in rats, mice, and humans ([Sweeney et al. 2012a](#); [Sweeney et al. 2012b](#)).
  - 2a. Are the conclusions reached based on EPA's evaluation of the models scientifically supported? Do the revised PBPK models adequately represent RDX toxicokinetics? Are the model assumptions and parameters clearly presented and scientifically supported? Are the uncertainties in the model appropriately considered and discussed?
  - 2b. The average concentration of RDX in arterial blood (expressed as area under the curve) was selected over peak concentration as the dose metric for interspecies extrapolation for oral points of departure (PODs) derived from rat data. Is the choice of dose metric for each hazard sufficiently explained and appropriate? The mouse PBPK model was not used to derive PODs for noncancer or cancer endpoints because of uncertainties in the model and because of uncertainties associated with selection of a dose metric for cancer endpoints. Is this decision scientifically supported?
  - 2c. In Section 2.1.3 of the draft assessment, an UF of 10 for human variation is applied in the derivation of the RfD. Does the toxicokinetic modeling support the use of a different factor instead?
3. **Hazard identification and dose-response assessment<sup>1</sup>.** In Chapter 1, the draft assessment evaluates the available human, animal, and mechanistic studies to identify health outcomes that may result from exposure to RDX. In Chapter 2, the draft assessment develops organ/system- specific reference values for the health outcomes identified in Chapter 1, then selects overall reference values for each route of exposure. The draft assessment uses EPA's guidance documents (see <http://www.epa.gov/iris/basic-information-about-integrated-risk-information-system#guidance>) to reach the following conclusions.

---

<sup>1</sup> [Note: As suggested by the Chemical Assessment Advisory Committee panel that reviewed the draft IRIS assessment of benzo[a]pyrene, the charge questions in this section are organized by health outcome, with a question on each hazard identification followed by questions on the corresponding organ/system-specific toxicity values. This suggestion, however, entails some redundancy, as some questions apply equally to multiple health outcomes.]

3a. Nervous system effects

- i. **Nervous system hazard** (Sections 1.2.1, 1.3.1). The draft assessment concludes that nervous system toxicity is a human hazard of RDX exposure. Please comment on whether the available human, animal, and mechanistic studies support this conclusion. Are all hazards to the nervous system adequately assessed? Is there an appropriate endpoint to address the spectrum of effects?
- ii. **Nervous system-specific toxicity values** (Section 2.1.1). Please comment on whether the selection of studies reporting nervous system effects is scientifically supported and clearly described. Considering the difference in toxicokinetics between gavage and dietary administration (described in Appendix C, Section C.1, and in the context of specific hazards in the toxicological review), is it appropriate to consider the [Crouse et al. \(2006\)](#) study, which used gavage administration? Is the characterization of convulsions as a severe endpoint, and the potential relationship to mortality, appropriately described?
- iii. **Points of departure for nervous system endpoints** (Section 2.1.2). Is the selection of convulsions as the endpoint to represent this hazard scientifically supported and clearly described? Are the calculations of PODs for these studies scientifically supported and clearly described? Is the calculation of the HEDs for these studies scientifically supported and clearly described? Does the severity of convulsions warrant the use of a benchmark response level of 1% extra risk? Is calculation of the lower bound on the benchmark dose (BMDL) for convulsions appropriate and consistent with the EPA's Benchmark Dose Guidance?
- iv. **Uncertainty factors for nervous system endpoints** (Section 2.1.3). Is the application of uncertainty factors to these PODs scientifically supported and clearly described? The subchronic and database uncertainty factors incorporate multiple considerations; please comment specifically on the scientific rationale for the application of a subchronic uncertainty factor of 1 and a UF<sub>D</sub> of 3<sup>2</sup>.
- v. **Nervous system-specific reference dose** (Section 2.1.4). Is the organ/system-specific reference dose derived for nervous system effects scientifically supported and clearly characterized?

---

<sup>2</sup> Note that the UF<sub>D</sub> applies to each of the hazards identified in the toxicological review.

3b. Kidney and other urogenital system effects

- i. **Kidney and other urogenital system hazard** (Sections 1.2.2, 1.3.1). The draft assessment concludes that kidney and other urogenital system toxicity is a potential human hazard of RDX exposure. Please comment on whether the available human, animal, and mechanistic studies support this conclusion. Are all hazards to kidney and urogenital system adequately assessed? Is the selection of suppurative prostatitis as the endpoint to represent this hazard scientifically supported and clearly described?
- ii. **Kidney and other urogenital system-specific toxicity values** (Section 2.1.1). Is the selection of the [Levine et al. \(1983\)](#) study that describes kidney and other urogenital system effects scientifically supported and clearly described?
- iii. **Points of departure for kidney and other urogenital system endpoints** (Section 2.1.2). Is the calculation of a POD for this study scientifically supported and clearly described? Is the calculation of the HED for this study scientifically supported and clearly described?
- iv. **Uncertainty factors for kidney and other urogenital system endpoints** (Section 2.1.3). Is the application of uncertainty factors to the POD scientifically supported and clearly described?
- v. **Kidney and other urogenital system-specific reference dose** (Section 2.1.4). Is the organ/system-specific reference dose derived for kidney and other urogenital system effects scientifically supported and clearly characterized?

3c. Developmental and reproductive system effects

- i. **Developmental and reproductive system hazard** (Sections 1.2.3, 1.3.1). The draft assessment concludes that there is suggestive evidence of male reproductive effects associated with RDX exposure, based on evidence of testicular degeneration in male mice. The draft assessment did not draw any conclusions as to whether developmental effects are a human hazard of RDX exposure. Please comment on whether the available human, animal, and mechanistic studies support these decisions. Are other hazards to human reproductive and developmental outcome adequately addressed?
- ii. **Reproductive system-specific toxicity values** (Section 2.1.1). Is the selection of the [Lish et al. \(1984\)](#) study that describes male reproductive system effects scientifically supported and clearly described?
- iii. **Points of departure for reproductive system endpoints** (Section 2.1.2). Is the calculation of a POD for this study scientifically supported and clearly described? Is the calculation of the HED for this study scientifically supported and clearly described?

- iv. **Uncertainty factors for reproductive system endpoints** (Section 2.1.3). Is the application of uncertainty factors to the POD scientifically supported and clearly described?
- v. **Reproductive system-specific reference dose** (Section 2.1.4). Is the organ/system-specific reference dose derived for reproductive system effects scientifically supported and clearly characterized?
- 3d. **Other noncancer hazards** (Sections 1.2.4, 1.2.6, 1.3.1). The draft assessment did not draw any conclusions as to whether liver, ocular, musculoskeletal, cardiovascular, immune, or gastrointestinal effects are human hazards of RDX exposure. Please comment on whether the available human, animal, and mechanistic studies support this decision. Are other non-cancer hazard adequately described?
- 3e. **Cancer**
- i. **Cancer hazard** (Sections 1.2.5, 1.3.2). There are plausible scientific arguments for more than one hazard descriptor as discussed in Section 1.3.2. The draft assessment concludes that there is *suggestive evidence of carcinogenic potential* for RDX, and that this descriptor applies to all routes of human exposure. Please comment on whether the available human, animal, and mechanistic studies support these conclusions.
- ii. **Cancer-specific toxicity values** (Section 2.3.1). As noted in EPA's 2005 *Guidelines for Carcinogen Risk Assessment*, "When there is suggestive evidence, the Agency generally would not attempt a dose-response assessment, as the nature of the data generally would not support one; however, when the evidence includes a well-conducted study, quantitative analyses may be useful for some purposes, for example, providing a sense of the magnitude and uncertainty of potential risks, ranking potential hazards, or setting research priorities." Does the draft assessment adequately explain the rationale for quantitative analysis, considering the uncertainty in the data and the suggestive nature of the weight of evidence, and is the selection of the [Lish et al. \(1984\)](#) study for this purpose scientifically supported and clearly described?
- iii. **Points of departure for cancer endpoints** (Section 2.3.2, 2.3.3). Are the calculations of PODs and oral slope factors scientifically supported and clearly described?
4. **Dose-response analysis.** In Chapter 2, the draft assessment uses the available human, animal, and mechanistic studies to derive candidate toxicity values for each hazard that is credibly associated with RDX exposure in Chapter 1, identify an organ/system-specific RfD, then selects an overall toxicity value for each route of exposure. The draft assessment uses EPA's guidance documents (see <http://www.epa.gov/iris/basic-information-about-integrated-risk-information-system#guidance>) in the following analyses.

1  
2 **4a. Oral reference dose for effects other than cancer** (Sections 2.1.5–2.1.8). The draft  
3 assessment presents an overall oral reference dose of  $3 \times 10$  mg/kg-day, based on nervous  
4 system effects as described in the [Crouse et al. \(2006\)](#) study. Is this selection scientifically  
5 supported and clearly described, including consideration of mortality as described in  
6 Section 2.1.6, and consideration of the organ/system-specific reference dose derived from  
7 the toxicity study by [Cholakakis et al. \(1980\)](#) that is lower (by approximately fivefold) as  
8 described in Section 2.1.4?  
9

10 **4b. Inhalation reference concentration for effects other than cancer** (Section 2.2).  
11 The draft assessment does not derive an inhalation reference concentration as the  
12 available studies were insufficient to characterize inhalation hazard and conduct dose-  
13 response analysis, and no toxicokinetic studies of RDX were available to support  
14 development of a PBPK inhalation model. If you believe that the available data might  
15 support an inhalation reference concentration, please describe how one might be derived.  
16

17 **4c. Oral slope factor for cancer** (Section 2.3.3–2.3.4). The draft assessment presents an  
18 overall oral slope factor of 0.038 per mg/kg-day based on the combination of liver and  
19 lung tumors in female mice. Is this derivation scientifically supported and clearly  
20 described?  
21

22 **4d. Inhalation unit risk for cancer** (Section 2.4). The draft assessment does not derive an  
23 inhalation unit risk because inhalation carcinogenicity data were not available, nor were  
24 toxicokinetic studies of inhalation of RDX available to support development of an  
25 inhalation PBPK model. If you believe that the available data might support an inhalation  
26 unit risk, please describe how one might be derived.  
27

- 28 5. **Executive Summary.** Does the executive summary clearly and adequately present the major  
29 conclusions of the assessment?  
30

## References

- Cholakakis, JM; Wong, LCK; Van Goethem, DL; Minor, J; Short, R; Sprinz, H; Ellis, HV, III. (1980). Mammalian toxicological evaluation of RDX. (DAMD17-78-C-8027). Kansas City, MO: Midwest Research Institute.
- Crouse, LCB; Michie, MW; Major, M; Johnson, MS; Lee, RB; Paulus, HI. (2006). Subchronic oral toxicity of RDX in rats. (Toxicology Study No. 85-XC-5131-03). Aberdeen Proving Ground, MD: U.S. Army Center for Health Promotion and Preventive Medicine.
- Levine, BS; Lish, PM; Furedi, EM; Rac, VS; Sagartz, JM. (1983). Determination of the chronic mammalian toxicological effects of RDX (twenty-four month chronic toxicity/carcinogenicity study of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) in the Fischer 344 rat): Final report--phase V. Chicago, IL: IIT Research Institute.
- Lish, PM; Levine, BS; Furedi, EM; Sagartz, JM; Rac, VS. (1984). Determination of the chronic mammalian toxicological effects of RDX: Twenty-four month chronic toxicity/carcinogenicity study of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) in the B6C3F1 hybrid mouse (Volumes 1-3). (ADA181766. DAMD17-79-C-9161). Fort Detrick, Frederick, MD: U.S. Army Medical Research and Development Command. <http://oai.dtic.mil/oai/oai?verb=getRecord&metadataPrefix=html&identifier=ADA181766>
- Sweeney, LM; Gut, CP, Jr; Gargas, ML; Reddy, G; Williams, LR; Johnson, MS. (2012a). Assessing the non-cancer risk for RDX (hexahydro-1,3,5-trinitro-1,3,5-triazine) using physiologically based pharmacokinetic (PBPK) modeling [Review]. Regul Toxicol Pharmacol 62: 107-114. <http://dx.doi.org/10.1016/j.yrtph.2011.12.007>
- Sweeney, LM; Okolica, MR; Gut, CP, Jr; Gargas, ML. (2012b). Cancer mode of action, weight of evidence, and proposed cancer reference value for hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX). Regul Toxicol Pharmacol 64: 205-224. <http://dx.doi.org/10.1016/j.yrtph.2012.07.005>



## APPENDIX B: EDITORIAL COMMENTS

### **Specific comments on text and presentation of data on reproductive and developmental toxicity**

Page 1-39, line 30: Reference to historic controls should be deleted. It is not valid to compare with historical controls (Ward et al. 1979) because it was done by different investigators and no quantitative level of what constitutes testicular degeneration is presented.

Page 1-39, line 33: Add "and a 14% decrease in testis weight" It is useful to add this because this was the only study in which testis weight showed a corresponding decrease with germ cell degeneration, as would be expected.

Page 1-40, line 4: Delete text and references about increases in testis weights. The only significant increase was for relative testis weight and this was really due to a loss of body weight.

Page 1-40, lines 8-21: The presentation and discussion of the results of the two-generation and dominant lethal studies should be combined. They were not separate studies; in fact the same male rats were used for both. Moreover, they show that same result: decrease in yield of pregnancies from males exposed to 50 mg/kg-day. The only difference was the treatment of the females.

### **Specific comments on presentation of data (Tables, Figures) on reproductive and developmental toxicity**

1. The Tables and Figures are well planned and show the important features that need to be presented. However SAB concludes that Table 1 of this report should be added as it compares all the rodent studies in one table, facilitating comparisons showing support and discrepancy.

2. In Table 1-9, the relative testes weights should be deleted. Relative testis weight is affected by changes in body weights, which in our experience does not have effects on testis weights of adult animals. Absolute testis weights are a better measure of testicular toxicity of an agent. The relative testis weights just clutter up the table and add little information on the toxicity of RDX.

3. Table 1-9, Page 1-42: In the presentation of the data of Levine et al. (1983), the data on "SDMS" (spontaneous death or moribund sacrifice) rats should be deleted. Their significance is open to question and they aren't given much weight in the discussion.

4. Table 1-9, Page 1-44: The data on incidence of germ degeneration of Levine et al. (1981a,b, 1990) at 12 and 15 mg/kg-day should be deleted. These were observed on dead rats (all rats in these groups died). Incidentally the numbers were reversed: the value for 1/10 was for the 12 mg/kg-day dose and 1/9 was for 15 mg/kg-day.

5. Table 1-9 (footnote, Page 1-44) Also reference to historic controls for comparison of testicular degeneration reported by Lish et al. (1984) should be deleted.

6. The testis weight data from Cholakakis et al. (1980) (Table 1-9, last entry on Page 1-43) on F2 weanlings does not belong in the male reproductive effect section. It is not indicative of direct effect on testis weight and there is no follow-up to determine whether or not adult testis weights will be affected. Rather it belongs in the developmental effects section (Table 1-10).

7. Figure 1-3 could provide a useful comparison of doses from various studies. However, to achieve maximum impact, the data should be grouped as follows: mouse; rat 2-year chronic; rat 13-week subchronic. The study using gavage should be noted since the effective dose seems to be dependent on method of oral administration. Footnote (1), indicating that the non-significant



1 change in testis weight in Hart et al. (1976) was a slight increase, was confusing and should be  
2 deleted; anyway that is covered by the symbol that there was no significant change.

3 8. Figure 1-3. Additionally, it may be a matter of rote procedure, but the decision to  
4 highlight only statistically significant findings in the exposure-response array is deceptive  
5 because the two studies identified with statistical significant findings (Levine 1990 and 1983)  
6 were not considered meaningful results, but the nonstatistically significant finding in Lish et al.  
7 (1984) is the finding for male reproductive effects that is debated heavily in this document and is  
8 not highlighted. Perhaps add an explanation via footnotes.

9 9. Table 1-10 Page 1-46: Reconsider the use of term 'offspring survival' to categorize  
10 'prenatal mortality' as offspring survival is more commonly associated with postnatal outcome.

11 10. Figure1-4: typo in spelling of 'significantly' in the key.  
12

## 1 **APPENDIX C: SUGGESTIONS ON THE FORMAT FOR EPA's CHARGE QUESTIONS**

2  
3 The CAAC-RDX panel has the following observations on the charge questions based on  
4 experience during the review meeting:  
5

61) Charge questions on the calculation of points of departure for organ/system-specific reference  
7 dose did not account for the possibility that the panel may not agree with the selection of the  
8 specific endpoint for derivation of a POD (as is the case for the use of suppurative prostatitis  
9 for derivation of a POD for kidney and other urogenital system effects, and the use of testicular  
10 degeneration for the derivation of a POD for male reproductive effect).  
11

### 12 ***Suggestions***

- 13 • There should be a question if the panel agrees with the selection of a specific endpoint for  
14 derivation of a POD, before the question if the calculation of the POD is scientifically  
15 supported and clearly described.
- 16 • There should also be a question on whether there is an alternative approach.